


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1991

[Full Table of Content](#)



91/R/CS/212 Seasonal Effects of Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (1992) *91/R/CS/212 Seasonal Effects of Take-all - W. Wheat* ; Yields Of The Field Experiments 1991, pp 55 - 56 - DOI: <https://doi.org/10.23637/ERADOC-1-46>

91/R/CS/212

SEASONAL EFFECTS OF TAKE-ALL

Object: To study the incidence of take-all (*Gaeumannomyces graminis*) in continuous w. wheat and in first, second and third w. wheats after a break - Great Harpenden I.

Sponsors: D. Hornby, R.J. Gutteridge.

The 14th year, w. wheat.

For previous years see 78-90/R/CS/212.

Design: 3 randomised blocks of 8 plots.

Whole plot dimensions: 5.33 x 10.0.

Treatments:

PREVCROP Previous crops before w. wheat 1991:

	78	79	80	81	82	83	84	85	86	87	88	89	90
W13	W	W	W	W	W	W	W	W	W	W	W	W	W
BE2 W4	W	BE	W	W	BE	W	W	BE	BE	W	W	W	W
BE1 W4	W	W	W	W	W	W	W	W	BE	W	W	W	W
BE1 W6	BE	W	W	BE	W	W	BE	W	W	W	W	W	W
BE1 W7	W	W	BE	W	W	BE	W	W	W	W	W	W	W
BE1 W2	W	BE	W	W	BE	W	W	BE	W	W	BE	W	W
BE1 W1	W	W	BE	W	W	BE	W	W	BE	W	W	BE	W
BE1	BE	W	W	BE	W	W	BE	W	W	BE	W	W	BE

BE = s. beans, W = w. wheat

Basal applications: Manures: (0:16:36) at 980 kg. 'Nitram' at 410 kg. Weedkillers: Isoproturon at 1.3 kg and pendimethalin at 1.3 kg with the insecticide in 200 l. Glyphosate at 0.72 kg with a wetting agent, 'Frigate' at 1.0 l, in 200 l. Fungicides: Fenpropimorph at 0.38 kg in 200 l. Chlorothalonil at 0.49 kg and flutriafol at 0.078 kg with fenpropimorph at 0.38 kg in 200 l. Insecticide: Deltamethrin at 6.2 g.

Seed: Mercia, sown at 170 kg.

Cultivations, etc.: - PK applied: 3 Sept, 1990. Ploughed: 10 Sept. Cultivated by rotary grubber: 26 Sept. Rotary harrowed, seed sown: 27 Sept. Isoproturon, pendimethalin and the insecticide applied: 15 Nov. N applied: 3 Apr, 1991. Fenpropimorph alone applied: 10 May. Chlorothalonil, flutriafol and fenpropimorph applied: 20 June. Glyphosate with wetting agent applied: 13 Aug. Combine harvested: 20 Aug.

NOTE: Plant and soil samples were taken frequently during the season to assess take-all. Additional soil samples were taken to measure the suppressiveness of the soil to the take-all fungus.

91/R/CS/212

GRAIN TONNES/HECTARE

***** Tables of means *****

PREVCROP	
W13	6.91
BE2 W4	6.97
BE1 W4	7.21
BE1 W6	7.08
BE1 W7	7.41
BE1 W2	6.83
BE1 W1	6.51
BE1	7.72
Mean	7.08

*** Standard errors of differences of means ***

PREVCROP
0.191

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	14	0.234	3.3
GRAIN MEAN DM%	87.6		
PLOT AREA HARVESTED	0.00216		