


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1991

[Full Table of Content](#)



91/R/CS/323 Cereal Sequence and Take-all - W. Wheat, W. Barley, S. Barley, W. Triticale, W. Oats

Rothamsted Research

Rothamsted Research (1992) *91/R/CS/323 Cereal Sequence and Take-all - W. Wheat, W. Barley, S. Barley, W. Triticale, W. Oats* ; Yields Of The Field Experiments 1991, pp 67 - 69 - DOI: <https://doi.org/10.23637/ERADOC-1-46>

91/R/CS/323

CEREAL SEQUENCES AND TAKE-ALL

Object: To study the effects on take-all (*Gaeumannomyces graminis*) and yield of including triticale in cereal sequences - West Barnfield II.

Sponsors: R.J. Gutteridge, D. Hornby, R.D. Prew.

The fourth year, w. barley, w. oats, w. triticale, w. wheat, s. barley.

For previous years see 88-90/R/CS/323

Design: 3 randomised blocks of 26 plots.

Whole plot dimensions: 3.0 x 10.0.

CROPSEQ Crop sequences (1988, 1989, 1990 and 1991 respectively):

W W S S	W. wheat, w. wheat, s. barley, s. barley
B B B B	W. barley, w. barley, w. barley, w. barley
O B B B	W. oats, w. barley, w. barley, w. barley
B O B B	W. barley, w. oats, w. barley, w. barley
B B O B	W. barley, w. barley, w. oats, w. barley
S B S B	S. barley, w. barley, s. barley, w. barley
T T B B	W. triticale, w. triticale, w. barley, w. barley
T B T B	W. triticale, w. barley, w. triticale, w. barley
W B W B	W. wheat, w. barley, w. wheat, w. barley
W W B B	W. wheat, w. wheat, w. barley, w. barley
B B B O	W. barley, w. barley, w. barley, w. oats
T T T O	W. triticale, w. triticale, w. triticale, w. oats
W W W O	W. wheat, w. wheat, w. wheat, w. oats
T T T T	W. triticale, w. triticale, w. triticale, w. triticale
B B T T	W. barley, w. barley, w. triticale, w. triticale
O T T T	W. oats, w. triticale, w. triticale, w. triticale
T O T T	W. triticale, w. oats, w. triticale, w. triticale
T T O T	W. triticale, w. triticale, w. oats, w. triticale
W W T T	W. wheat, w. wheat, w. triticale, w. triticale
W T W T	W. wheat, w. triticale, w. wheat, w. triticale
W W W W	W. wheat, w. wheat, w. wheat w. wheat
B B W W	W. barley, w. barley, w. wheat, w. wheat
O W W W	W. oats, w. wheat, w. wheat, w. wheat
W O W W	W. wheat, w. oats, w. wheat, w. wheat
W W O W	W. wheat, w. wheat, w. oats, w. wheat
T T W W	W. triticale, w. triticale, w. wheat, w. wheat

Standard applications:

W. cereals: Manures: (0:16:36) at 300 kg. N at 30 kg to all w. cereals followed by N at 150 kg (w. barley), 170 kg (w. wheat), and 120 kg (w. oats and w. triticale), all as 'Nitram'.
Weedkillers: Isoxaben at 76 g and methabenzthiazuron at 1.6 kg in 200 l. Fluroxypyr at 0.15 kg in 200 l. Fungicides: Carbendazim at 0.15 kg and prochloraz at 0.40 kg with tridemorph at 0.26 kg in 200 l.

91/R/CS/323

Standard applications:

S. barley: Manures: (0:16:36) at 300 kg. N at 120 kg as 'Nitram'.
Weedkiller: Metsulfuron-methyl at 6.0 g with fenpropimorph in
200 l. Fungicides: Carbendazim at 0.15 kg and prochloraz at
0.40 kg with tridemorph at 0.26 kg in 200 l. Fenpropimorph at
0.75 kg.

Seed: W. barley: Magie, sown at 130 kg.
W. oats: Image, sown at 130 kg.
W. triticale: Lasko, sown at 140 kg.
W. wheat: Mercia, sown at 170 kg.
S. barley: Klaxon, sown at 120 kg.

Cultivations, etc.:-

W. cereals: Ploughed, furrow pressed: 4 Sept, 1990. PK applied:
14 Sept. Rotary harrowed twice: 17 Sept. Rotary harrowed, seed
sown: 19 Sept. Isoxaben and methabenzthiazuron applied: 20 Sept.
First N applied: 4 Mar, 1991. Fluroxypyr applied: 12 Apr. Second
N applied: 15 Apr. Fungicides applied: 23 Apr. Combine
harvested: 12 Aug (w. barley), 20 Aug (w. wheat), 27 Aug (w. oats
and w. triticale).

S. barley: Ploughed, furrow pressed: 4 Sept, 1990. PK applied:
14 Sept. Rotary harrowed twice: 17 Sept. N applied: 14 Mar, 1991.
Rotary harrowed twice, seed sown: 15 Mar. Carbendazim, prochloraz
and tridemorph applied: 23 Apr. Weedkiller with fenpropimorph
applied: 11 June. Combine harvested: 15 Aug.

NOTE: Plants were sampled in April, June and July to assess take-all,
eyespot and sharp eyespot. Soil cores were taken after harvest to
assess take-all infectivity.

91/R/CS/323

GRAIN TONNES/HECTARE

***** Tables of means *****

CROPSEQ	
W W S S	5.40
B B B B	5.89
O B B B	7.22
B O B B	7.14
B B O B	7.36
S B S B	7.16
T T B B	6.63
T B T B	5.79
W B W B	6.54
W W B B	5.71
B B B O	5.98
T T T O	6.35
W W W O	5.08
T T T T	4.19
B B T T	4.51
O T T T	4.38
T O T T	3.93
T T O T	5.54
W W T T	4.21
W T W T	4.42
W W W W	5.65
B B W W	6.34
O W W W	7.38
W O W W	7.05
W W O W	7.05
T T W W	5.04
Mean	5.84

*** Standard errors of differences of means ***

CROPSEQ
0.389

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	50	0.476	8.2
GRAIN MEAN DM%	84.9		
PLOT AREA HARVESTED	0.00226		