


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1992

[Full Table of Content](#)



92/R/BS/2 Spray Timings and Bydv- S. Barley

Rothamsted Research

Rothamsted Research (1993) *92/R/BS/2 Spray Timings and Bydv- S. Barley* ; Yields Of The Field Experiments 1992, pp 118 - 120 - DOI: <https://doi.org/10.23637/ERADOC-1-47>

92/R/BS/2

SPRING BARLEY

SPRAY TIMINGS AND BYDV

Object: To investigate the optimum strategy for controlling barley yellow dwarf virus (BYDV) in spring barley in relation to sowing date, aphid immigration and subsequent population development - Little Hoos.

Sponsors: J. Mann, N. Carter, R.T. Plumb.

Design: 3 randomised blocks of 16 plots with external dummy plots and arranged to allow estimation of the effects of neighbouring plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments:

S P DATE	Dates of sowing and of applying pirimicarb:
E 0	Sown March 1992, no pirimicarb
E D1	" " pirimicarb applied after emergence
E D2	" " " " 10 days after emergence
E D3	" " " " 20 " " "
E D1 D2	" " " " after emergence and 10 days later
E D1 D3	" " " " after emergence and 20 days later
E D2 D3	" " " " 10 days and 20 days after emergence
E D1D2D3	" " " " after emergence, 10 days and 20 days later
L 0	Sown April, no pirimicarb
L D2	" " pirimicarb applied after emergence
L D3	" " " " 10 days after emergence
L D4	" " " " 20 " " "
L D2 D3	" " " " after emergence and 10 days later
L D2 D4	" " " " after emergence and 20 days later
L D3 D4	" " " " 10 days and 20 days after emergence
L D2D3D4	" " " " after emergence, 10 days and 20 days later

Experimental diary:

- 30-Aug-91 : B : Straw chopped.
- 30-Oct-91 : B : Ploughed.
- 17-Mar-92 : T : S P DATE E 0, E D1, E D2, E D3, E D1 D2, E D1 D3, E D2 D3, E D1D2D3: Rotary harrowed. Alexis, dressed Cerevax, drilled at 160 kg.
- 19-Mar-92 : B : Rolled.
- 22-Apr-92 : B : 34.5% N at 440 kg.
- 22-Apr-92 : T : S P DATE L 0, L D2, L D3, L D4, L D2 D3, L D2 D4, L D3 D4, L D2D3D4: Spring-tine cultivated, rotary harrowed, Alexis, dressed Cerevax, drilled at 160 kg.

92/R/BS/2

Experimental diary:

24-Apr-92 : B : Rolled.
05-May-92 : T : **S P DATE** E D1, E D1 D2, E D1 D3, E D1D2D3: Aphox at
0.28 kg in 300 l.
15-May-92 : T : **S P DATE** E D2, E D1 D2, E D2 D3, E D1D2D3, L D2,
L D2 D3, L D2 D4, L D2D3D4: Aphox at 0.28 kg in
300 l.
19-May-92 : B : Duplosan New System CMPP at 2.0 l, Vindex at 1.0 l and
Calixin at 0.70 l in 200 l.
22-May-92 : T : **S P DATE** E D3, E D1 D3, E D2 D3, E D1D2D3, L D3,
L D2 D3, L D3 D4, L D2D3D4: Aphox at 0.28 kg in
300 l.
04-Jun-92 : T : **S P DATE** L D4, L D2 D4, L D3 D4, L D2D3D4: Aphox at
0.28 kg in 300 l.
09-Jun-92 : B : Radar at 0.50 l in 200 l.
18-Aug-92 : B : Combine harvested.

Previous crops: W. oilseed rape 1990, w. wheat 1991.

- NOTES:** (1) Plant samples were taken to assess aphid numbers in spring and summer.
(2) Symptoms of BYDV were visually assessed and leaf samples taken for enzyme-linked immunosorbent assay in spring and summer.
(3) Number of ears was counted before harvest and thousand grain weights determined at harvest.

92/R/BS/2

GRAIN TONNES/HECTARE

***** Tables of means *****

S P DATE	
E 0	7.93
E D1	7.89
E D2	8.23
E D3	8.10
E D1 D2	7.89
E D1 D3	8.12
E D2 D3	8.13
E D1D2D3	7.92
L 0	5.50
L D2	5.48
L D3	5.51
L D4	5.42
L D2 D3	5.78
L D2 D4	5.62
L D3 D4	5.62
L D2D3D4	5.72
Mean	6.80

*** Standard errors of differences of means ***

S P DATE
0.162

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	30	0.198	2.9
GRAIN MEAN DM%	85.0		
PLOT AREA HARVESTED	0.00230		