


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1992

[Full Table of Content](#)



### 92/R/RAW/6 Light Leaf Spot Study - W. Oilseed Rape

#### Rothamsted Research

Rothamsted Research (1993) *92/R/RAW/6 Light Leaf Spot Study - W. Oilseed Rape* ; Yields Of The Field Experiments 1992, pp 131 - 132 - DOI: <https://doi.org/10.23637/ERADOC-1-47>

92/R/RAW/6

WINTER OILSEED RAPE

LIGHT LEAF SPOT STUDY

**Object:** To study methods for the prediction and detection of early infection of light leaf spot (*Pyrenopeziza brassicae*) and subsequent effects on crop phenology and yield - Great Knott III.

**Sponsors:** L. Figueroa, B.D.L. Fitt.

**Design:** 4 randomised blocks of 6 plots split into 2 sub plots.

**Whole plot dimensions:** 9.0 x 21.0.

**Treatments:**

Whole plots

1. **FUNGCIDE** Time of fungicide application:

NONE	None
PROCAUT	Prochloraz in autumn
PROCWINT	Prochloraz in winter
PROCSPR	Prochloraz in spring
IPROSUM	Iprodione in summer
PA+PS+IS	Prochloraz in autumn and spring, iprodione in summer

Sub plots

2. **VARIETY** Varieties:

CAPRCORN	Capricorn
FALCON	Falcon

**Experimental diary:**

28-Aug-91 : B : Straw chopped.  
05-Sep-91 : B : Ploughed, rolled.  
06-Sep-91 : T : **VARIETY** CAPRCORN: Rotary harrowed. Capricorn, dressed Lindex-Plus FS, drilled at 9.7 kg.  
06-Sep-91 : T : **VARIETY** FALCON: Rotary harrowed. Falcon, dressed Lindex-Plus FS, drilled at 8.0 kg.  
06-Sep-91 : B : Rolled.  
13-Oct-91 : B : Pilot at 0.15 l with Cropspray 11 E at 2.5 l in 200 l.  
11-Nov-91 : B : Benazalox at 0.75 kg and Butisan S at 1.5 l in 200 l.  
02-Dec-91 : T : **FUNGCIDE** PROCAUT, PA+PS+IS: Sportak 45 at 0.55 l in 200 l.  
17-Feb-92 : T : **FUNGCIDE** PROCWINT: Sportak 45 at 0.55 l in 200 l.  
19-Feb-92 : B : 34.5% N at 220 kg.  
21-Feb-92 : B : Benazalox at 1.5 kg in 200 l.  
25-Mar-92 : B : 34.5% N at 290 kg.  
09-Apr-92 : T : **FUNGCIDE** PROCSPR, PA+PS+IS: Sportak 45 at 1.1 l in 200 l.  
04-Jun-92 : T : **FUNGCIDE** IPROSUM, PA+PS+IS: Rovral Flo at 2.0 l in 200 l.

92/R/RAW/6

**Experimental diary:**

15-Jul-92 : B : Reglone at 3.0 l with Farmon Blue at 0.26 l in 260 l.  
 20-Jul-92 : B : Combine harvested.

Previous crops: W. wheat 1990 and 1991.

**NOTE:** Disease incidence on the crop was assessed at intervals during the growing season. Air-borne inoculation was monitored from autumn to spring using bait plants and a Burkard spore sampler. At harvest oil and glucosinolate levels in the seed were determined.

**GRAIN (AT 90% DRY MATTER) TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

FUNGCIDE VARIETY	NONE	PROCAUT	PROCWINT	PROCSRP	IPROSUM	PA+PS+IS	Mean
CAPRCORN	4.60	4.55	4.38	4.67	4.45	4.53	4.53
FALCON	3.75	3.91	3.68	3.96	3.68	4.06	3.84
Mean	4.18	4.23	4.03	4.31	4.07	4.29	4.18

\*\*\* Standard errors of differences of means \*\*\*

	FUNGCIDE	VARIETY	FUNGCIDE VARIETY
	0.220	0.077	0.257
Except when comparing means with the same level(s) of FUNGCIDE			0.188

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	15	0.311	7.4
BLOCK.WP.SP	18	0.266	6.4

GRAIN MEAN DM% 79.3

SUB PLOT AREA HARVESTED 0.00483