

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1992

[Full Table of Content](#)



92/R/LN1 Varieties and Disease - Linseed

Rothamsted Research

Rothamsted Research (1993) *92/R/LN1 Varieties and Disease - Linseed* ; Yields Of The Field Experiments 1992, pp 155 - 156 - DOI: <https://doi.org/10.23637/ERADOC-1-47>

92/R/LN/1

LINSEED

VARIETIES AND DISEASE

Object: To investigate the effects of pathogens on varieties of linseed - White Horse II.

Sponsors: B.D.L. Fitt, J. Harold.

Design: 4 randomised blocks of 10 plots.

Whole plot dimensions: 3.0 x 15.0.

Treatments: All combinations of:-

1. **VARIETY** Varieties:

ANTARES
BARBARA
MCGREGOR
NORLIN
CD 1747

2. **FUNGICIDE** Fungicides:

NONE None
APPLIED Prochloraz seed dressing, iprodione during flowering,
prochloraz and carbendazim plus maneb post-
flowering

Experimental diary:

16-Oct-91 : B : Straw burnt.
28-Nov-91 : B : Farmyard manure at 10 t.
04-Dec-91 : B : Ploughed.
18-Mar-92 : B : PK as (0:18:36) at 940 kg.
10-Apr-92 : B : 34.5% N at 220 kg.
11-Apr-92 : B : Spring-tine cultivated.
21-Apr-92 : T : **VARIETY:** All varieties drilled at 550 seeds per square
metre.
: T : **FUNGICIDE APPLIED:** Seed, dressed Prelude 20LF, drilled.
22-Apr-92 : B : Rolled.
02-Jun-92 : B : Basagran at 2.0 l and Vindex at 1.0 l in 200 l.
24-Jun-92 : T : **FUNGICIDE APPLIED:** Rovral Flo at 2.0 l in 200 l.
08-Jul-92 : T : **FUNGICIDE APPLIED:** Sportak 45 at 0.90 l in 200 l.
23-Jul-92 : T : **FUNGICIDE APPLIED:** Delsene M Flowable at 5.0 l in 200 l.
05-Sep-92 : B : Challenge at 3.0 l in 400 l.
29-Sep-92 : B : Combine harvested.

Previous crops: W. wheat 1990, linseed 1991.

NOTE: Emergence counts were made in spring. Samples were taken fortnightly from June to September for disease assessment. Oil content of the grain was measured.

92/R/LN/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

FUNGCIDE VARIETY	NONE	APPLIED	Mean
ANTARES	2.41	2.55	2.48
BARBARA	1.74	2.12	1.93
MCGREGOR	2.31	2.42	2.36
NORLIN	2.01	2.30	2.16
CD 1747	2.15	2.32	2.23
Mean	2.12	2.34	2.23

*** Standard errors of differences of means ***

VARIETY	FUNGCIDE	VARIETY FUNGCIDE
0.087	0.055	0.123

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	27	0.174	7.8
GRAIN MEAN DM%	88.7		
PLOT AREA HARVESTED	0.00276		