

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1992

[Full Table of Content](#)



---

### Linseed

#### Rothamsted Research

Rothamsted Research (1993) *Linseed* ; Yields Of The Field Experiments 1992, pp 155 - 158 - **DOI:** <https://doi.org/10.23637/ERADOC-1-47>

92/R/LN/1

LINSEED

VARIETIES AND DISEASE

**Object:** To investigate the effects of pathogens on varieties of linseed - White Horse II.

**Sponsors:** B.D.L. Fitt, J. Harold.

**Design:** 4 randomised blocks of 10 plots.

**Whole plot dimensions:** 3.0 x 15.0.

**Treatments:** All combinations of:-

1. **VARIETY** Varieties:

ANTARES  
BARBARA  
MCGREGOR  
NORLIN  
CD 1747

2. **FUNGICIDE** Fungicides:

NONE None  
APPLIED Prochloraz seed dressing, iprodione during flowering,  
prochloraz and carbendazim plus maneb post-  
flowering

**Experimental diary:**

16-Oct-91 : B : Straw burnt.  
28-Nov-91 : B : Farmyard manure at 10 t.  
04-Dec-91 : B : Ploughed.  
18-Mar-92 : B : PK as (0:18:36) at 940 kg.  
10-Apr-92 : B : 34.5% N at 220 kg.  
11-Apr-92 : B : Spring-tine cultivated.  
21-Apr-92 : T : **VARIETY:** All varieties drilled at 550 seeds per square  
metre.  
: T : **FUNGICIDE APPLIED:** Seed, dressed Prelude 20LF, drilled.  
22-Apr-92 : B : Rolled.  
02-Jun-92 : B : Basagran at 2.0 l and Vindex at 1.0 l in 200 l.  
24-Jun-92 : T : **FUNGICIDE APPLIED:** Rovral Flo at 2.0 l in 200 l.  
08-Jul-92 : T : **FUNGICIDE APPLIED:** Sportak 45 at 0.90 l in 200 l.  
23-Jul-92 : T : **FUNGICIDE APPLIED:** Delsene M Flowable at 5.0 l in 200 l.  
05-Sep-92 : B : Challenge at 3.0 l in 400 l.  
29-Sep-92 : B : Combine harvested.

Previous crops: W. wheat 1990, linseed 1991.

**NOTE:** Emergence counts were made in spring. Samples were taken fortnightly from June to September for disease assessment. Oil content of the grain was measured.

92/R/LN/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

FUNGCIDE VARIETY	NONE	APPLIED	Mean
ANTARES	2.41	2.55	2.48
BARBARA	1.74	2.12	1.93
MCGREGOR	2.31	2.42	2.36
NORLIN	2.01	2.30	2.16
CD 1747	2.15	2.32	2.23
Mean	2.12	2.34	2.23

\*\*\* Standard errors of differences of means \*\*\*

VARIETY	FUNGCIDE	VARIETY FUNGCIDE
0.087	0.055	0.123

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	27	0.174	7.8
GRAIN MEAN DM%	88.7		
PLOT AREA HARVESTED	0.00276		

92/R/LN/2

LINSEED

WEED COMPETITION AND LINSEED

**Object:** To investigate the effects of weed competition on the growth and yield of linseed - Fosters Corner.

**Sponsor:** P.J.W. Lutman.

**Design:** 3 randomised blocks of 10 plots.

**Whole plot dimensions:** 3.0 x 10.0.

**Treatments:** All combinations of:-

1. **SPECIES** Weed species:
- |          |   |
|----------|---|
| CUL OATS | Cultivated oats ( <i>Avena sativa</i> ) to simulate wild oats ( <i>A. fatua</i> ) |
| CHICKWD  | Chickweed ( <i>Stellaria media</i> )  |
2. **DENSITY** Density of established plants (number of plants per square metre):
- |    | Cultivated oats | Chickweed |
|----|-----------------|-----------|
| D0 | 0               | 0         |
| D1 | 6               | 57        |
| D2 | 24              | 159       |
| D3 | 49              | 182       |
| D4 | 119             | 465       |

**NOTES:** (1) Target plant populations (number of plants per square metre):

	D0	D1	D2	D3	D4
Cultivated oats	0	10	50	100	200
Chickweed	0	100	200	400	800

(2) Weed species were sown in the central 2 m of each plot.

**Experimental diary:**

- 06-Sep-91 : B : Straw chopped.  
06-Dec-91 : B : Ploughed (start).  
02-Jan-92 : B : Ploughed (finish).  
22-Apr-92 : B : 34.5% N at 220 kg.  
          : B : Spring-tine cultivated.  
23-Apr-92 : B : Harrowed.  
23-Apr-92 : T : **SPECIES** CUL OATS: Dula drilled.  
          : T : **SPECIES** CHICKWD: Broadcast by hand.  
23-Apr-92 : B : Rotary harrowed, Antares, dressed Prelude 20LF, drilled at 48 kg, harrowed.  
24-Apr-92 : B : Rolled.  
09-Jun-92 : T : **SPECIES** CUL OATS: Basagran at 2.0 l and Vindex at 1.0 l in 220 l.  
08-Sep-92 : T : Hand harvested (start).  
09-Sep-92 : T : Hand harvested (finish).

92/R/LN/2

Previous crops: W. wheat 1990, s. beans 1991.

**NOTE:** Weed counts were made and dry weight samples were taken at intervals during the growing season.

**GRAIN (AT 90% DRY MATTER) TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

DENSITY SPECIES	D0	D1	D2	D3	D4	Mean
CUL OATS	1.62	1.34	0.98	0.65	0.47	1.01
CHICKWD	1.63	1.27	1.39	1.43	1.43	1.43
Mean	1.63	1.31	1.18	1.04	0.95	1.22

\*\*\* Standard errors of differences of means \*\*\*

SPECIES	DENSITY	SPECIES
		DENSITY
	0.065	0.144
	0.102	

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	18	0.177	14.5

GRAIN MEAN DM% \*

PLOT AREA HARVESTED 0.00020