

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1992

[Full Table of Content](#)



### 92/W/CS/347 Green Crops for Set-aside - Leys, Tumbledown, W. Oats

#### Rothamsted Research

Rothamsted Research (1993) 92/W/CS/347 *Green Crops for Set-aside - Leys, Tumbledown, W. Oats* ;  
Yields Of The Field Experiments 1992, pp 76 - 78 - DOI: <https://doi.org/10.23637/ERADOC-1-47>

92/W/CS/347

**GREEN CROPS FOR SET-ASIDE**

**Object:** To obtain information on the establishment and maintenance of sown crops and unsown vegetation in a long-term, up to five-year, set-aside area given no chemical inputs. Effects on soil nitrate and leaching after ploughing are also studied - Woburn, Horsepool Lane Close II.

**Sponsors:** R.D. Prew, E.T.G. Bacon, M.V. Hewitt, D.P. Yeoman.

**Design:** 6 randomised blocks of 6 plots.

**Whole plot dimensions:** 6.5 x 26.0.

The third year, w. oats, ryegrass, clover, tumbledown.

For previous years see 90-91/W/CS/347.

**Treatments:**

<b>CROPS</b>	Crops, cumulative since 1990:
RY LF	Ryegrass, cuttings left in situ
RY+CL LF	Ryegrass + clover, cuttings left in situ
RY+CL RE	Ryegrass + clover, cuttings removed
RY+N RE	Ryegrass given 100 kg N in spring, cuttings removed
TU LF	Tumbledown, natural regrowth, cuttings left in situ
W OATS	Winter oats

**NOTES:** (1) Yields were taken from the w. oats and from the ley plots, from which cuttings were removed.

(2) Ryegrass and clover were sown in 1989.

**Experimental diary:**

- 18-Sep-91 : T : **CROPS** W OATS: Ploughed.
- 21-Sep-91 : T : **CROPS** W OATS: Rolled.
- 25-Sep-91 : T : **CROPS** W OATS: Rotary cultivated, Image drilled at 152 kg rolled.
- 20-Mar-92 : T : **CROPS** RY+CL RE: 21 kg P<sub>2</sub>O<sub>5</sub> and 84 kg K<sub>2</sub>O to balance PK in crop removed last year.  
RY+N RE: 53 kg P<sub>2</sub>O<sub>5</sub> and 264 kg K<sub>2</sub>O to balance PK in crop removed last year.
- 10-Apr-92 : T : **CROPS** W OATS: 34.5% N at 290 kg.
- 15-May-92 : T : **CROPS** W OATS: Dorin at 1.0 l and Ally at 30 g in 300 l.
- 27-May-92 : T : **CROPS** RY LF, RY+CL LF, RY+CL RE and RY+N RE: Cut.
- 28-May-92 : T : **CROPS** TU LF: Cut.
- 02-Jun-92 : T : **CROPS** RY+CL RE and RY+N RE: Cuttings removed.
- 15-Jul-92 : T : **CROPS** RY LF, RY+CL LF, RY+CL RE, RY+N RE and TU LF: Cut.
- 15-Jul-92 : T : **CROPS** RY+CL RE and RY+N RE: Cuttings removed.
- 31-Jul-92 : T : **CROPS** W OATS: Combine harvested.
- 15-Sep-92 : T : **CROPS** RY LF, RY+CL LF, RY+CL RE and RY+N RE: Cut.
- 16-Sep-92 : T : **CROPS** TU LF: Cut.
- 16-Sep-92 : T : **CROPS** RY+CL RE and RY+N RE: Cuttings removed.

92/W/CS/347

**NOTE:** Soil nitrogen was measured in autumn 1991 and spring 1992. Ground cover, plant numbers, plant height and growth stages were estimated in spring and autumn 1992. Soil seedbank samples were taken in autumn 1992.

**GRASS**

**1ST CUT (27/5/92) DRY MATTER TONNES/HECTARE**

\*\*\*\*\* Tables of means\*\*\*\*\*

CROPS	RY+CL RE	RY+N RE	Mean
	1.44	4.27	2.85

\*\*\* Standard errors of differences of means \*\*\*

**CROPS**  
0.280

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	5	0.484	17.0

1ST CUT MEAN DM% 23.3

**2ND CUT (15/7/92) DRY MATTER TONNES/HECTARE**

\*\*\*\*\* Tables of means\*\*\*\*\*

CROPS	RY+CL RE	RY+N RE	Mean
	1.22	0.37	0.79

\*\*\* Standard errors of differences of means \*\*\*

**CROPS**  
0.116

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	5	0.201	25.3

2ND CUT MEAN DM% 23.3

92/W/CS/347

**GRASS**

**3RD CUT (15/9/92) DRY MATTER TONNES/HECTARE**

\*\*\*\*\* Tables of means\*\*\*\*\*

<b>CROPS</b>	<b>RY+CL RE</b>	<b>RY+N RE</b>	<b>Mean</b>
	2.02	0.75	1.38

\*\*\* Standard errors of differences of means \*\*\*

**CROPS**  
0.170

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

<b>Stratum</b>	<b>d.f.</b>	<b>s.e.</b>	<b>cv%</b>
BLOCK.WP	5	0.295	21.3

3RD CUT MEAN DM% 27.4

**TOTAL OF 3 CUTS DRY MATTER TONNES/HECTARE**

\*\*\*\*\* Tables of means\*\*\*\*\*

<b>CROPS</b>	<b>RY+CL RE</b>	<b>RY+N RE</b>	<b>Mean</b>
	4.67	5.38	5.03

\*\*\* Standard errors of differences of means \*\*\*

**CROPS**  
0.478

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

<b>Stratum</b>	<b>d.f.</b>	<b>s.e.</b>	<b>cv%</b>
BLOCK.WP	5	0.828	16.5

TOTAL OF 3 CUTS MEAN DM% 24.7

PLOT AREA HARVESTED 0.00264

**W. OATS**

**GRAIN TONNES/HECTARE** 4.65

**GRAIN MEAN DM%** 86.5

**PLOT AREA HARVESTED** 0.00572