

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1992

[Full Table of Content](#)



---

### 92/W/CS/356 Set-aside Study - W. Wheat, Fallow

#### Rothamsted Research

Rothamsted Research (1993) *92/W/CS/356 Set-aside Study - W. Wheat, Fallow* ; Yields Of The Field Experiments 1992, pp 83 - 84 - DOI: <https://doi.org/10.23637/ERADOC-1-47>

92/W/CS/356

**SET-ASIDE STUDY**

**Object:** To compare different treatments of land temporarily withdrawn from arable cropping and to study their effects on nitrate leaching and on subsequent wheat crops - Woburn, Horsepool Lane Close I.

**Sponsors:** R.D. Prew, E.T.G. Bacon, M.V. Hewitt, D.P. Yeoman, J.F. Jenkyn, R.J. Gutteridge, W. Powell, J. Ashby.

**Associate sponsors:** D.L.O. Smith, I. Shield, M.D. Helps.

The second year, w. wheat.

For first year see 91/W/CS/356.

**Design:** 3 randomised blocks of 7 plots split into 8.

**Whole plot dimensions:** 10.0 x 24.0.

**Treatments:** All combinations of:-

Whole plot

1. **LAND TRT[91]** Land treatment in 1991, after w. wheat 1990 (all treatments ploughed autumn 1991 before w. wheat):
  - CA WW Cultivated in autumn, sown to w. wheat
  - CA RA Cultivated in autumn, sown to ryegrass in autumn, topped in spring
  - SA CA FA Straw chopped and spread in autumn, cultivated in autumn, sown to forage rape in autumn, topped in spring
  - CA CS Cultivated in autumn, cultivated in spring
  - SA CS Straw chopped and spread in autumn, cultivated in spring
  - WT Weeds topped
  - WT CS TS Weeds topped, cultivated in spring, trefoil sown in spring, topped

Sub plots

2. **N** Nitrogen fertilizer (kg N) as 'Nitro-Chalk':
  - 0
  - 80
  - 120
  - 160
  - 200
  - 240
  - 280

**NOTE:** An additional fallow sub plot was present, systematically arranged on one side of each whole plot.

92/W/CS/356

**Experimental diary.**

W. wheat:

- 18-Sep-91 : B : Ploughed.
- 25-Sep-91 : T : Rolled.
- 02-Oct-91 : T : Rotary cultivated, Mercia drilled at 140 kg.
- 18-Mar-92 : T : Folimat at 1.12 l in 200 l.
- 01-Apr-92 : T : Duplosan New System CMPP at 2.0 l and Asset at 2.0 l in 200 l.
- 09-Apr-92 : T : N 80, 120, 160, 200, 240 and 280: Applied as 27% N.
- 15-May-92 : T : Dorin at 1.0 l and Ally at 0.03 kg in 300 l.
- 08-Jun-92 : T : Halo at 2.0 l in 200 l.
- 29-Jul-92 : T : Barclay Gallup at 2.0 l with Team at 0.30 l in 300 l.
- 01-Sep-92 : T : Combine harvested.

Fallow:

- 18-Sep-91 : B : Ploughed.
- 25-Apr-92 : T : Rotary cultivated.

**NOTE:** Amounts of soil nitrogen and plant dry matter were measured in autumn and spring. Assessment of plant cover was made in spring. Diseases were assessed in spring and summer.

**GRAIN TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

	N	0	80	120	160	200	240	280	Mean
<b>LAND TRT[91]</b>									
CA WW	5.37	6.65	6.24	6.25	6.21	7.27	6.46	6.35	
CA RA	4.26	6.53	6.86	6.91	7.29	7.42	7.47	6.68	
SA CA FA	5.08	6.19	7.35	7.31	7.39	6.35	6.76	6.63	
CA CS	7.62	7.56	7.61	7.08	7.30	6.58	6.46	7.17	
SA CS	6.45	7.90	7.50	7.64	7.27	7.75	7.14	7.38	
WT	5.88	7.16	6.74	7.35	7.43	7.69	7.70	7.13	
WT CS TS	7.56	7.18	7.39	7.30	6.62	6.34	5.16	6.79	
Mean	6.03	7.02	7.10	7.12	7.07	7.06	6.74	6.88	

\*\*\* Standard errors of differences of means \*\*\*

LAND TRT[91]	N	LAND TRT[91]
		N
	0.262	0.273
Except when comparing means with the same level(s) of		0.718
<b>LAND TRT[91]</b>		0.722

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	12	0.321	4.7
BLOCK.WP.SP	84	0.884	12.9

GRAIN MEAN DM% 83.0      SUB PLOT AREA HARVESTED 0.00199