Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1992



Full Table of Content

# 92/W/CS/375 Take-all Epidemics - W. Wheat

# **Rothamsted Research**

Rothamsted Research (1993) 92/W/CS/375 Take-all Epidemics - W. Wheat; Yields Of The Field Experiments 1992, pp 90 - 91 - DOI: https://doi.org/10.23637/ERADOC-1-47

### 92/W/CS/375

# TAKE-ALL EPIDEMICS

**Object:** To determine whether severe take-all (Gaeumannomyces graminis) can be caused by artificial inoculum in winter wheat and to determine the distribution of such infection within the crop - Woburn, Butt Close I.

Sponsors: G. L. Bateman, D. Hornby.

The first year, w. wheat

Design: 3 randomised blocks of 6 x 2, plus 2 extra plots.

Whole plot dimensions: 6.0 x 2.50.

Treatments: All combinations of:-

SOW DATE Date of sowing:

E Early

L Late (4 weeks later)

2. INOCULTN Weight (kg) of inoculated oat seed applied by combine drill in autumn and spring:

	Autumn	(E)	Autumn	(L)	Spring
0	Nil		Nil		-
1	Nil		200		-
2	200		200		_
3	400		200		_
30	400		200		Nil
35	400		200		500

plus 2 extra plots, systematically arranged with treatments 0 and 2:

NOTE: INOCULTN 0, 1, 30: Nil occurs where empty drill drawn across plots.

#### Experimental diary:

19-Sep-91 : B : Sub-soiled and ploughed.

23-Sep-91 : B : Rotary harrowed.

: T : INOCULTN 0 and 1: Empty drill drawn across plots.
INOCULTN 2, 3, 30 and 3S: Inoculated oat seed applied by combine drill.

: T : SOW DATE E: Mercia, undressed, drilled at 180 kg.

14-Oct-91 : T : SOW DATE L: Rotary harrowed.

: T : INOCULTN 0: Empty drill drawn across plots.

INOCULTN 1, 2, 3, 30 and 3S: Inoculated oat seed applied
by combine drill.

: T : SOW DATE L: Mercia, undressed, drilled at 180 kg.

06-Jan-92 : B : PK as (0:16:36) at 740 kg.

31-Jan-92 : B : Stomp 400 at 2.5 1 and Stefes IPU at 1.0 1 in 200 1.

10-Mar-92 : B : 34.5% N at 116 kg.

25-Mar-92 : T : INOCULTN 30: Empty drill drawn across plots.

INOCULTN 3S: Inoculated oat seed applied by combine
 drill.

#### 92/W/CS/375

# Experimental diary:

22-Apr-92 : B : 34.5% N at 464 kg.

09-Jun-92 : B : Impact Excel at 2.0 1 in 200 1.

29-Aug-92 : B : Combine harvested.

Previous crops: Potatoes 1990, navy beans 1991.

NOTE: Plant samples were taken for take-all assessment in autumn, spring and summer. Extra plots were sampled throughout the growing season to measure take-all and plant growth in relation to growth stage.

#### GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

INOCULTN	0	1	2	3	30	35	Mean
SOW DATE							
E	4.27	2.77	2.81	2.50	1.38	2.61	2.72
L	3.75	2.40	2.08	1.04	1.24	1.63	2.03
Mean	4.01	2.59	2.45	1.77	1.31	2.12	2.37

\*\*\* Standard errors of differences of means \*\*\*

SOW DATE	INOCULTN	SOW DATE
		INOCULTN
0.342	0.592	0.837

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*

Stratum	d.f.	s.e.	CV%
BLOCK.WP	22	1.025	43.2

GRAIN MEAN DM% 81.7

PLOT AREA HARVESTED 0.00132