


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1993

[Full Table of Content](#)



93/R/RAW/2 Varieties and Fungicides - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1994) *93/R/RAW/2 Varieties and Fungicides - W. Oilseed Rape* ; Yields Of The Field Experiments 1993, pp 124 - 125 - DOI: <https://doi.org/10.23637/ERADOC-1-48>

93/R/RAW/2

WINTER OILSEED RAPE

VARIETIES AND FUNGICIDES

Object: To investigate the effects of fungicides on a range of low glucosinolate varieties - Bones Close.

Sponsors: V.J. Church, B.D.L. Fitt.

Design: 4 randomised blocks of 2 x 6 plots.

Whole plot dimensions: 3.0 x 21.0.

Treatments: All combinations of:-

1. **VARIETY** Varieties:

CAPRCORN	Capricorn
ENVOL	Envol
FALCON	Falcon
LIBRAVO	Libravo
SAMOURAI	Samourai
ROCKET	Rocket

2. **FUNGICIDE** Fungicides:

NONE	None
PR+CA+IP	Prochloraz and carbendazim in autumn and spring, iprodione in summer

Experimental diary:

27-Jul-92 : B : Shallow cultivated with Bomford Dynadrive.
28-Jul-92 : B : Rolled.
18-Aug-92 : B : Sting CT at 2.0 l in 200 l.
20-Aug-92 : B : Ploughed, furrow pressed.
26-Aug-92 : T : **VARIETY** CAPRCORN, ENVOL, FALCON, LIBRAVO, SAMOURAI, ROCKET: Rotary harrowed. All varieties dressed Lindex-Plus FS, drilled at 120 seeds per square metre.
22-Oct-92 : B : Benazalox at 0.75 l and Butisan S at 1.5 l in 200 l.
08-Dec-92 : T : **FUNGICIDE** PR+CA+IP: Sportak 45 at 1.1 l and Tripart Defensor FL at 0.50 l in 200 l.
18-Feb-93 : B : 34.5% N at 170 kg.
08-Mar-93 : T : **FUNGICIDE** PR+CA+IP: Sportak 45 at 1.1 l and Tripart Defensor FL at 0.50 l in 200 l.
23-Mar-93 : B : 34.5% N at 370 kg.
18-May-93 : T : **FUNGICIDE** PR+CA+IP: Rovral Flo at 2.0 l in 200 l.
19-Jul-93 : B : Stefes Diquat at 3.0 l with Vassgro Spreader at 0.40 l in 400 l.
26-Jul-93 : B : Combine harvested.

Previous crops: W. barley 1991 and 1992.

93/R/RAW/2

NOTE: Samples were taken throughout the year for disease assessments on leaves, stems and pods. Oil content of seed was measured after harvest.

GRAIN (AT 90% DM) TONNES/HECTARE

***** Tables of means *****

FUNGCIDE VARIETY	NONE	PR+CA+IP	Mean
CAPRCORN	4.17	4.68	4.42
ENVOL	4.59	5.08	4.84
FALCON	4.08	4.27	4.18
LIBRAVO	4.16	4.68	4.42
SAMOURAI	4.33	4.73	4.53
ROCKET	3.87	3.99	3.93
Mean	4.20	4.57	4.39

*** Standard errors of differences of means ***

VARIETY	FUNGCIDE	VARIETY FUNGCIDE
0.090	0.052	0.127

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	33	0.180	4.1
GRAIN MEAN DM%	83.2		
PLOT AREA HARVESTED	0.00483		