


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1993

[Full Table of Content](#)



### 93/W/RAS/1 Weed Competition and N in Spring Rape - S. Oilseed Rape

#### Rothamsted Research

Rothamsted Research (1994) *93/W/RAS/1 Weed Competition and N in Spring Rape - S. Oilseed Rape* ; Yields Of The Field Experiments 1993, pp 142 - 143 - DOI:  
<https://doi.org/10.23637/ERADOC-1-48>

93/W/RAS/1

SPRING OILSEED RAPE

WEED COMPETITION AND N IN SPRING RAPE

**Object:** To investigate the effect of nitrogen fertilizer on the competitiveness of charlock and chickweed in spring rape - Woburn, Butt Furlong.

**Sponsor:** P.J.W. Lutman.

**Design:** 3 randomised blocks of 3 x 4 plots.

**Whole plot dimensions:** 3.0 x 10.0.

**Treatments:** All combinations of:-

1. **WEED SP** Weed species sown:

S ARVEN	<i>Sinapsis arvensis</i> (charlock)
S MEDIA	<i>Stellaria media</i> (chickweed)
NONE	None

2. **N** Rates of nitrogen (kg N):

50  
100  
150  
200

**Experimental diary:**

04-Mar-93 : B : Ploughed  
05-Mar-93 : B : Rolled.  
29-Mar-93 : T : N 50, 100, 150, 200: Applied.  
          : T : WEED SP S ARVEN, S MEDIA: Weed seeds broadcast.  
          : B : Rotary harrowed.  
          : B : Starlight, dressed Lindex-Plus FS, drilled at 150 seeds per square metre.  
04-Jun-93 : B : Decis at 0.50 l in 200 l.  
13-Aug-93 : T : N 50, 100: Hand harvested.  
19-Aug-93 : T : N 150, 200: Hand harvested.

**NOTES:** (1) Weed seeds were sown in the central 2m strip along the length of the plot.  
(2) Target plant populations (plants per square metre) and sowing rates (g per square metre) were respectively S ARVEN 100, 4.0, S MEDIA 400, 0.8.  
(3) Population counts of crop and weeds were made in May. Ground cover was assessed throughout the season and plant biomass samples were taken in June and August.  
(4) The WEED SP S ARVEN plots were not harvested because the weed density was too high, no rape survived. One plot was severely grazed by rabbits. The yield of this plot with treatment combination WEED SP NONE N 50 was lost. An estimated value was used in the analysis.

93/W/RAS/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

	N	50	100	150	200	Mean
<b>WEED SP</b>						
S MEDIA		1.05	1.68	1.78	1.29	1.45
NONE		1.40	1.37	2.73	2.06	1.89
Mean		1.22	1.53	2.25	1.67	1.67

\*\*\* Standard errors of differences of means \*\*\*

WEED SP	N	WEED SP
		N
0.244	0.345	0.488

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	13	0.598	35.8

GRAIN MEAN DM% 90.2

PLOT AREA HARVESTED 0.00010