

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1993

[Full Table of Content](#)



---

### 93/R/LN/1 Varieties and Disease - Linseed

#### Rothamsted Research

Rothamsted Research (1994) *93/R/LN/1 Varieties and Disease - Linseed* ; Yields Of The Field Experiments 1993, pp 159 - 160 - DOI: <https://doi.org/10.23637/ERADOC-1-48>

93/R/LN/1

LINSEED

VARIETIES AND DISEASE

**Object:** To investigate the effects of pathogens on varieties of linseed - Sawyers II.

**Sponsors:** B.D.L. Fitt, J. Harold.

**Design:** 4 randomised blocks of 2 X 5 plots.

**Whole plot dimensions:** 3.0 x 15.0.

**Treatments:** All combinations of:-

1. **VARIETY** Varieties:

ANTARES  
BARBARA  
MCGREGOR  
NORLIN  
CD 1747

2. **FUNGICIDE** Fungicidal sprays:

NONE None  
APPLIED Iprodione during flowering, prochloraz and carbendazim plus maneb post-flowering

**NOTE:** Basal fungicidal seed dressing (prochloraz) was applied.

**Experimental diary:**

22-Jan-93 : B : Ploughed.  
19-Apr-93 : B : Heavy spring-tine cultivated.  
20-Apr-93 : T : **VARIETY** ANTARES, BARBARA, MCGREGOR, NORLIN, CD 1747:  
Rotary harrowed, varieties, dressed Prelude 20LF,  
drilled at 700 seeds per square metre.  
12-May-93 : B : 34.5% N at 220 kg.  
13-May-93 : B : Decis at 0.30 l in 200 l.  
01-Jun-93 : B : Ally at 30 g in 200 l.  
28-Jun-93 : T : **FUNGICIDE** APPLIED: Rovral Flo at 2.0 l in 200 l.  
22-Jul-93 : T : **FUNGICIDE** APPLIED: Sportak 45 at 0.90 l in 200 l.  
: T : **FUNGICIDE** APPLIED: Tripart Defensor FL at 1.0 l in 200 l.  
04-Aug-93 : T : **FUNGICIDE** APPLIED: Multi-W FL at 5.0 l in 200 l.  
06-Sep-93 : B : Stefes Diquat at 3.0 l with Vassgro Spreader at 0.30 l  
in 260 l.  
23-Oct-93 : T : Combine harvested.

Previous crops: W. wheat 1991, s. wheat 1992.

93/R/LN/1

**NOTE:** Emergence counts were made. Samples taken fortnightly between June and October for visual assessment of disease and assessment of pathogenic fungi by isolation on agar. Plant populations prior to harvest and oil content of seed were measured.

**GRAIN (AT 90% DM) TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

FUNGCIDE VARIETY	NONE	APPLIED	Mean
ANTARES	2.03	2.16	2.10
BARBARA	2.41	2.86	2.63
MCGREGOR	2.71	2.80	2.76
NORLIN	2.30	2.34	2.32
CD 1747	2.38	2.75	2.56
Mean	2.37	2.58	2.47

\*\*\* Standard errors of differences of means \*\*\*

FUNGCIDE	VARIETY	FUNGCIDE VARIETY
0.070	0.111	0.157

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	27	0.222	9.0

GRAIN MEAN DM% 87.1

PLOT AREA HARVESTED 0.00276