Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1993



Full Table of Content

## 93/W/P/1 Double Cropping - Potatoes

### **Rothamsted Research**

Rothamsted Research (1994) 93/W/P/1 Double Cropping - Potatoes; Yields Of The Field Experiments 1993, pp 167 - 168 - DOI: https://doi.org/10.23637/ERADOC-1-48

#### 93/W/P/1

#### POTATOES

#### DOUBLE CROPPING

Object: To study the effects of growing two crops of potatoes in one season on the population dynamics of *Globodera pallida* - Woburn, Mill Dam Close III.

Sponsors: K. Evans, P.D. Halford.

Design: 3 blocks of 2 x 2 plots with external dummy plots.

Whole plot dimensions: 3.0 x 6.0.

Treatments: All combinations of:-

1. CROPPING Cropping:

DOUBLE Double crop: Pentland Javelin, then Costella

MAIN Main crop: Costella

NEMACIDE Nematicide applied:

NONE None

OXAMYL Oxamyl at 4.0 kg

#### Experimental diary:

10-Aug-92 : B : Subsoiled to 45 cm with times 1.5 m apart

21-Jan-93 : B : Ploughed.

25-Jan-93 : B : Covered with polythene sheeting.

24-Feb-93: T: CROPPING DOUBLE: Spike rotavated, PK as (0:24:24) at 1265 kg, 27% N at 666 kg.

: T : CROPPING DOUBLE, NEMACIDE OXAMYL: Oxamyl at 4.0 kg.

: T : CROPPING DOUBLE: Spike rotavated, Pentland Javelin, dressed Rizolex Flowable, planted. Rotalin at 3.5 1

in 250 1.

: B : Re-covered with polythene sheeting.

20-Apr-93 : T : CROPPING MAIN: PK as (0:24:24) at 1265 kg 27% N at 666 kg.

: T : CROPPING MAIN, NEMACIDE OXAMYL: Oxamyl at 4.0 kg.

: T : CROPPING MAIN: Spike rotavated, Costella planted.

22-Apr-93 : T : CROPPING DOUBLE: Re-covered with polythene sheeting.

12-May-93 : T : CROPPING MAIN: Arresin at 5.0 1 in 250 1.

14-May-93 : T : CROPPING DOUBLE: Removed polythene sheeting.

18-May-93 : T : CROPPING MAIN: Gramoxone 100 at 3.0 1 in 250 1.

17-Jun-93 : B : Chiltern Manex at 2.5 l with Intracrop BLA at 0.30 l in 300 l.

24-Jun-93 : T : CROPPING DOUBLE: Pentland Javelin lifted.

25-Jun-93: T: CROPPING DOUBLE: Haulm raked off, rotary cultivated, PK as (0:24:24) at 1265 kg, 27% N at 666 kg. Rotary

cultivated, ridged up, Costella planted.

08-Jul-93 : B : Irrigated 12.5 mm.

13-Jul-93 : B : Ashlade Mancozeb FL at 2.5 1 with Intracrop BLA at 0.30 1 in 300 1.

#### 93/W/P/1

#### Experimental diary:

30-Jul-93: B: Ashlade Mancozeb FL at 2.5 l with Intracrop BLA at 0.30 l and Polyverdol at 3.0 l in 300 l.

09-Aug-93 : T : CROPPING DOUBLE: Hand hoed.

13-Aug-93 : B : Ashlade Mancozeb FL at 2.5 1 with Intracrop BLA at 0.30 1 in 300 1.

24-Aug-93 : B : Ashlade Mancozeb FL at 2.5 1 with Intracrop BLA at 0.30 1 in 300 1.

01-Sep-93 : T : CROPPING MAIN: Costella lifted.

04-Sep-93: T: CROPPING DOUBLE: Brestan 60 at 0.5 kg in 300 l. 17-Sep-93: T: CROPPING DOUBLE: Brestan 60 at 0.5 kg in 300 l. 21-Oct-93: T: CROPPING DOUBLE: Costella lifted.

NOTES: (1) Potatoes were planted at 75 cm spacing between rows, 30 cm within rows.

- (2) Polyverdol is a foliar nutrient solution (8:8:6) containing chelated trace elements, vitamin B, and plant growth hormones.
- (3) Soil samples were taken before planting and after each crop for nematode counts. Root samples were taken six and ten weeks after planting each crop for assessment of nematode population densities. Disease assessments were made in spring and tubers were assessed for disease after each harvest.

#### TOTAL TUBERS TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

NEMACIDE	NONE	OXAMYL	Mean
CROPPING			
DOUBLE	17.7	18.6	18.1
MAIN	10.6	20.9	15.8
Mean	14.2	19.7	17.0

\*\*\* Standard errors of differences of means \*\*\*

CROPPING	NEMACIDE	CROPPING	
		NEMACIDE	
0.873	0.873	1.234	

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv*	
BLOCK.WP	6	1.511	8.9	

PLOT AREA HARVESTED \*