

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1993

[Full Table of Content](#)



93/R/CS/323 Cereal Sequence and Take-all - W. Barley, W. Oats, W. Triticale, W. Wheat

Rothamsted Research

Rothamsted Research (1994) *93/R/CS/323 Cereal Sequence and Take-all - W. Barley, W. Oats, W. Triticale, W. Wheat* ; Yields Of The Field Experiments 1993, pp 71 - 73 - DOI:

<https://doi.org/10.23637/ERADOC-1-48>

93/R/CS/323

CEREAL SEQUENCES AND TAKE-ALL

Object: To study the effects on take-all (*Gaeumannomyces graminis*) and yield of different cereals grown in various cereal sequences - West Barnfield II.

Sponsors: R.J. Gutteridge, D. Hornby, R.D. Prew.

The sixth year, w. barley, w. oats, w. triticale, w. wheat.

For previous years see 88-92/R/CS/323

Design: 3 randomised blocks of 26 plots.

Whole plot dimensions: 3.0 x 10.0.

CROPSEQ Crop sequences (1988, 1989, 1990, 1991, 1992 and 1993 respectively):

TTTTTT
OTTTOT
TOTTTO
TTOTTT
TTTOTT
WWWWWW
OWWWOW
WOWWOW
WWOWWW
WWWOWW
BBBBBB
OBBBOB
BOBBBO
BBOBBB
BBBOBB
WTWTWT
WBWBWB
TBTBTB
SBSBSB
WTTTIW
WWBBBW
TTBBBT
TIWWWT
BBWWWB
BTTTBT
WWSSSW

W = W. wheat
S = S. barley
B = W. barley
O = W. oats
T = W. triticale

93/R/CS/323

Experimental Diary:

05-Sep-92 : B : Scythe at 3.0 l in 200 l.
09-Sep-92 : B : Ploughed, to finish.
16-Sep-92 : B : Disced and rolled.
21-Sep-92 : B : PK as (0:18:36) at 300 kg..
28-Sep-92 : B : Sting CT at 2.0 l in 200 l.
01-Oct-92 : B : Re-ploughed.
06-Oct-92 : T : **CROPSEQ** Barley plots: Rotary harrowed, Magie, dressed
Cerevax, drilled at 350 seeds per square metre.
: T : **CROPSEQ** Oat plots: Rotary harrowed, Image, dressed
Ceresol, drilled at 350 seeds per square metre.
: T : **CROPSEQ** Triticale plots: Rotary harrowed, Lasko, dressed
Cerevax, drilled at 400 seeds per square metre.
: T : **CROPSEQ** Wheat plots: Rotary harrowed, Mercia, dressed
Cerevax, drilled at 380 seeds per square metre.
04-Mar-93 : B : 34.5%N at 90 kg.
15-Mar-93 : T : **CROPSEQ** Barley plots: Tigress at 2.5 l in 200 l.
15-Apr-93 : T : **CROPSEQ** Oats and triticale plots: 34.5% N at 368 kg.
: T : **CROPSEQ** Barley plots: 34.5% N at 435 kg.
: T : **CROPSEQ** Wheat plots: 34.5% N at 493 kg.
: T : **CROPSEQ** Wheat plots: Cheetah R at 2.0 l in 220 l.
: T : **CROPSEQ** Triticale plots: Hoegrass at 3.0 l in 220 l.
21-Apr-93 : B : Ally at 30 g and Starane 2 at 1.0 l in 200 l.
18-May-93 : B : Calirus at 2.0 l and Corbel at 0.50 l in 200 l.
08-Jun-93 : T : **CROPSEQ** Wheat plots: Halo at 2.0 l and Mistral at 0.50 l
in 200 l.
09-Jun-93 : T : **CROPSEQ** Oat plots: Mistral at 1.0 l in 200 l.
02-Aug-93 : T : **CROPSEQ** Barley plots: Combine harvested.
10-Aug-93 : B : **CROPSEQ** Wheat, oats, triticale plots: Combine
harvested.

NOTE: Plant samples were taken in April, June and July for take-all and eyespot assessments. Soil cores were taken after harvest to assess take-all infectivity.

93/R/CS/323

GRAIN TONNES/HECTARE

***** Tables of means *****

CROPSEQ	
TTTTTT	5.39
OTTTOT	5.77
TOTTTO	7.30
TTOTTT	4.99
TTTOTT	5.42
WWWWWW	5.90
OWWWOW	8.16
WOWWOW	6.60
WWOWWW	5.96
WWWOWW	7.78
BBBBBB	4.21
OBBBBB	6.63
BOBBBB	5.97
BBOBBB	4.02
BBBOBB	5.58
WTWTWT	5.18
WBWBWB	5.76
TBTBTB	5.38
SBSBSB	4.82
WWTTIW	6.25
WWBBBW	6.07
TTBBBT	4.26
TTWWWT	3.92
BBWWWB	5.76
BBTTTB	5.78
WWSSSW	5.91
Mean	5.72

*** Standard errors of differences of means ***

CROPSEQ
0.661

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	50	0.810	14.2
GRAIN MEAN DM%	83.7		
PLOT AREA HARVESTED	0.00228		