Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1993



Full Table of Content

93/R/CS/354 Sowing Dates and Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (1994) *93/R/CS/354 Sowing Dates and Take-all - W. Wheat ;* Yields Of The Field Experiments 1993, pp 85 - 86 - DOI: https://doi.org/10.23637/ERADOC-1-48

93/R/CS/354

SOWING DATES AND TAKE-ALL

Object: To study the effects of sequences of sowing dates and volunteers on take-all (*Gaeumannomyces graminis*) and yield of winter wheat -Little Knott I.

Sponsors: R.J. Gutteridge, D. Hornby.

The third year, w. wheat.

For previous years see 91-92/R/CS/354

Design: 4 randomised blocks of 5 plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments:

SOW SEQ	Sequences of sowing date in 1991, 1992 and 1993 and level						
	of volunteers in 1992 and 1993:						
EEE	Early in 1991, 1992 and 1993						
ELL	Early in 1991, late in 1992 and 1993						
E L+ L+	Early in 1991, late in 1992 and 1993, volunteers encouraged						
in second and third years							
LEE	Late in 1991, early in 1992 and 1993						
L L* L*	Late in 1991, 1992 and 1993, volunteers controlled in						
second and third years							

NOTE: On E L+ L+ volunteers simulated by sowing 50 kg wheat seed after cultivations on 15 Sept.

Experimental diary:

02-Sep-92 : B : Ploughed and furrow pressed. 15-Sep-92 : B : Rotary harrowed. : T : SOW SEQ E E E, L E E: Rotary harrowed, Mercia, dressed Cerevax, drilled at 380 seeds per square metre. 14-Oct-92 : T : SOW SEQ L L* L*: Rotary harrowed to control volunteers. : T : SOW SEQ: E L L, E L+ L+, L L* L*: Rotary harrowed, Mercia, dressed Cerevax, drilled at 380 seeds per square metre. 11-Mar-93 : B : Hytane 500 FW at 3.0 l and Stomp 400 at 2.5 l in 200 l. 15-Mar-93 : B : PK as (0:18:36) at 1250 kg. 20-Apr-93 : B : 34.5% N at 460 kg. 03-Jun-93 : B : Cheetah R at 2.5 l and Calixin at 0.70 l in 200 l. 08-Jun-93 : B : Halo at 2.0 l in 200 l. 17-Aug-93 : B : Combine harvested.

NOTE: Plant samples were taken in April and July for take-all assessment. Soil cores were taken after harvest to assess take-all infectivity.

This work is licensed under a <u>Creative Commons Attribution 4.0 International License</u>.

***** Table	es of means	****				
SOW SEQ	EEE	ELL	E L+ L+	LEE	L L* L*	Mean
	7.93	7.54	7.05	7.49	7.75	7.55
*** Standar	rd errors of	differ	ences of m	eans ***		
	SEQ .335					
***** Strat	cum standard	l errors	and coeff.	icients of	variation	*****
Stratum		d.f		s.e.	CV\$	
BLOCK.WP		1	2	0.474	6.3	
GRAIN MEAN	DM% 87.1					

PLOT AREA HARVESTED 0.00228

93/R/CS/354

GRAIN TONNES/HECTARE