Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1994



Full Table of Content

94/R/RAW/3 Effects of Behaviour Modifying Chemicals - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1995) 94/R/RAW/3 Effects of Behaviour Modifying Chemicals - W. Oilseed Rape; Yields Of The Field Experiments 1994, pp 112 - 113 - DOI:

https://doi.org/10.23637/ERADOC-1-49

94/R/RAW/3

WINTER OILSEED RAPE

EFFECTS OF BEHAVIOUR MODIFYING CHEMICALS

Object: To study the effects of behaviour modifying chemicals in the field on the pests of w. oilseed rape - Appletree.

Sponsors: L.E. Smart, M.M. Blight.

Design: 5 x 5 quasi-complete Latin square.

Whole plot dimensions: 9.0 x 9.0.

Treatments:

| CHEMICAL | Type o | of | behaviour | modifying | chemical | released: |
|----------|--------|----|-----------|-----------|----------|-----------|
| - | None | | | | | |
| A | A | | | | | |
| В | В | | | | | |
| C | C | | | | | |
| D | D | | | | | |

NOTE: The behaviour modifying chemicals were mixtures of host plant volatiles in various combinations. They were released from point sources above the crop from October 1993 until mid-June 1994.

Experimental diary:

```
19-Jul-93: B: Cultivated by rotary grubber.
29-Jul-93: B: Deep tine cultivated with vibrating tines 60 cm apart,
45 cm deep.
02-Aug-93: B: Rolled.
16-Aug-93: B: PK as (0:18:36) at 1250 kg.
18-Aug-93: B: Ploughed and furrow pressed, rolled.
23-Sep-93: B: Rotary harrowed, Envol, dressed Lindex-Plus FS, drilled at 120 seeds per m².
```

07-Oct-93 : B : Draza at 5.5 kg. 08-Mar-94 : B : 34.5% N at 162 kg. 11-Apr-94 : B : 34.5% N at 346 kg. 08-Aug-94 : B : Combine harvested.

Previous crops: Potatoes 1992, set-aside 1993.

NOTE: Numbers of cabbage stem flea beetle larvae were assessed in plant samples taken in December and February. Numbers of pollen beetles and seed weevils were assessed weekly throughout the spring and early summer. Pod samples were taken in June to assess seed weevil damage.

94/R/RAW/3

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

CHEMICAL

3.12 A 3.26 3.22 В 3.18 C D 2.98

Mean 3.15

*** Standard errors of differences of means ***

CHEMICAL

0.128

***** Stratum standard errors and coefficients of variation *****

Stratum

d.f.

s.e.

CV%

ROW.COL

12 0.203 6.4

GRAIN MEAN DM% 90.4

PLOT AREA HARVESTED 0.00207