


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1994

[Full Table of Content](#)



94/R/RAW/8 Weed Competition - Rape Drilling Date and Chickweed - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1995) *94/R/RAW/8 Weed Competition - Rape Drilling Date and Chickweed - W. Oilseed Rape* ; Yields Of The Field Experiments 1994, pp 124 - 126 - DOI:

<https://doi.org/10.23637/ERADOC-1-49>

94/R/RAW/8

WINTER OILSEED RAPE

WEED COMPETITION - RAPE DRILLING DATE AND CHICKWEED

Object: To study the effect of chickweed (*Stellaria media*) on the growth and yield of w. rape sown on three different dates - Appletree.

Sponsor: P.J.W. Lutman.

Design: 3 randomised blocks of 3 plots split into 6 sub-plots.

Whole plot dimensions: 3.0 x 14.0.

Treatments:

Whole plots

1. **SOW DATE** Target sowing date of w. rape:

SE	Early, 20-25 Aug
SM	Middle, 5-10 Sept
SL	Late, 15-20 Sept

Sub-plots

2. **WEED DEN** Weed density, plants per m²:

	SE	SM	SL
D0	0	0	0
D1	35	54	39
D2	130	156	112
D3	304	414	342
D4	790	887	659
OS	0	0	0

NOTES: (1) No herbicide applied to D0 plots, herbicide applied to OS plots. OS plots (SE and SM only) were also hand weeded in spring.

(2) Target weed densities were D0 and OS 0, D1 50, D2 150, D3 450 and D4 900 plants per m².

Experimental diary:

19-Jul-93 : B : Cultivated with rotary grubber.

29-Jul-93 : B : Deep tine cultivated with vibrating tines 60 cm apart and 45 cm deep.

02-Aug-93 : B : Rolled.

16-Aug-93 : B : PK as (0:18:36) at 1250 kg.

18-Aug-93 : B : Ploughed and furrow pressed, rolled.

25-Aug-93 : T : **SOW DATE** SE: Rotary harrowed, chickweed broadcast by hand. Rotary harrowed, Falcon, dressed Lindex-Plus FS, drilled at 120 seeds per m².

07-Sep-93 : T : **SOW DATE** SM: Rotary harrowed, chickweed broadcast by hand. Rotary harrowed, Falcon, dressed Lindex-Plus FS, drilled at 120 seeds per m².

94/R/RAW/8

Experimental diary:

23-Sep-93 : T : SOW DATE SL: Rotary harrowed, chickweed broadcast by hand. Rotary harrowed, Falcon, dressed Lindex-Plus FS, drilled at 120 seeds per m².
07-Oct-93 : B : Draza at 5.5 kg.
17-Dec-93 : T : WEED DEN OS: Kerb Flo at 1.75 l in 220 l.
: T : SOW DATE SE, SM, WEED DEN D1-D4: Synchemicals Dalapon at 3.4 kg in 220 l.
31-Jan-94 : T : SOW DATE SL, WEED DEN D1-D4: Synchemicals Dalapon at 3.4 kg in 220 l.
08-Mar-94 : B : 34.5% N at 162 kg.
11-Apr-94 : B : 34.5% N at 346 kg.
19-May-94 : T : SOW DATE SL: Fastac at 200 ml in 200 l.
21-Jul-94 : B : Landgold Diquat at 3.0 l with Vassgro Spreader at 300 ml in 300 l.
27-Jul-94 : B : Combine harvested.

Previous crops: Potatoes 1992, set-aside 1993.

NOTE: (1) Crop and weed densities were assessed and samples were taken to measure plant growth in December and March.
(2) The yields on two plots were treated as missing because of severe lodging, with treatment combinations:
SOW DATE SM SM
WEED DEN D2 OS
Estimated values were used in the analysis.

94/R/RAW/8

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

WEED DEN SOW DATE	D0	D1	D2	D3	D4	OS	Mean
SE	2.98	3.81	3.74	3.72	3.55	4.12	3.65
SM	2.16	3.55	3.58	3.54	3.70	4.00	3.42
SL	4.10	3.33	3.32	3.27	3.17	4.06	3.54
Mean	3.08	3.56	3.55	3.51	3.47	4.06	3.54

*** Standard errors of differences of means ***

SOW DATE	WEED DEN	SOW DATE WEED DEN
0.286	0.160	0.382

Except when comparing means with the same level(s) of
SOW DATE 0.278

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	4	0.351	9.9
BLOCK.WP.SP	28	0.340	9.6

GRAIN MEAN DM% 89.6

SUB PLOT AREA HARVESTED 0.00290