


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1994

[Full Table of Content](#)



### 94/R/BES/3 *bean* Flower Colour and Pheromone - S. Beans

#### Rothamsted Research

Rothamsted Research (1995) *94/R/BES/3 bean Flower Colour and Pheromone - S. Beans* ; Yields Of The Field Experiments 1994, pp 139 - 140 - DOI: <https://doi.org/10.23637/ERADOC-1-49>

94/R/BES/3

SPRING BEANS

BEAN FLOWER COLOUR AND PHEROMONE

**Object:** To compare the incidence of *Sitona lineatus* in purple and white flowered beans with and without insecticide and pheromone - Little Hoos.

**Sponsors:** L.E. Smart, M.M. Blight, R.T. Glinwood.

**Design:** 6 x 6 quasi-complete Latin square.

**Whole plot dimensions:** 6.0 x 6.0.

**Treatments:**

**TREATMNT** Variety, insecticide and pheromone:

A-	Alfred
AI	Alfred with insecticide
AP	Alfred with pheromone
C-	Caspar
CI	Caspar with insecticide
CP	Caspar with pheromone

**Experimental diary:**

29-Nov-93 : B : Deep tine cultivated.  
21-Mar-94 : B : Scythe at 2.0 l in 200 l.  
26-Mar-94 : B : Oxytril CM at 1.5 l with Starane 2 at 1.0 l in 200 l.  
30-Mar-94 : B : Heavy spring-tine cultivated, spring-tine cultivated.  
          : T : **TREATMNT** A-, AI, AP: Alfred drilled at 60 seeds per m<sup>2</sup>.  
          : T : **TREATMNT** C-, CI, CP: Caspar drilled at 60 seeds per m<sup>2</sup>.  
12-May-94 : T : **TREATMNT** AI, CI: Decis at 7.5 g in 200 l.  
08-Jul-94 : B : Bombardier at 1.0 l with Rovral Flo at 1.5 l and  
                  Pirimicarb 50 DG at 0.28 kg in 200 l.  
18-Aug-94 : T : Combine harvested.

Previous crops : S. barley 1992, potatoes 1993.

**NOTES:** (1) From late April, pheromone was released from a point source, which was hung above the crop at the plot centre.  
(2) Assessments were made of weevil larval numbers in root nodules at the end of May and of damage to leaves by adult weevils in April and May. Pitfall traps were used to assess weevil populations during the season.

94/R/BES/3

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

TREATMNT

A-	2.27
AI	2.36
AP	2.11
C-	2.47
CI	2.16
CP	1.71

Mean 2.18

\*\*\* Standard errors of differences of means \*\*\*

TREATMNT

0.267

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
---------	------	------	-----

ROW.COL	20	0.463	21.2
---------	----	-------	------

GRAIN MEAN DM% 80.3

PLOT AREA HARVESTED 0.00276