Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1994



Full Table of Content

# 94/R/P/4 Effects of Silver Scurf and Black Dot - Potatoes

# **Rothamsted Research**

Rothamsted Research (1995) 94/R/P/4 Effects of Silver Scurf and Black Dot - Potatoes; Yields Of The Field Experiments 1994, pp 163 - 166 - DOI: https://doi.org/10.23637/ERADOC-1-49

# 94/R/P/4

### POTATOES

# EFFECTS OF SILVER SCURF AND BLACK DOT

Object: To investigate the effect of planting seed with and without silver scurf into soil with or without black dot inoculum on disease in the subsequent crop at two harvest dates and after storage - Long Hoos VI/VII 5.

Sponsor: P.J. Read.

Design: 3 randomised blocks of 2 x 2 x 2 x 2 plots.

Whole plot dimensions: 1.5 x 5.7.

Treatments: All combinations of:-

## 1. VARIETY

ES Estima KE King Edward

2. SIL SCRF Infection with silver scurf (Helminthosporium solani):

S- None SD Infected

3. BLK DOT Infection with black dot (Colletotrichum coccodes):

B- None BD Infected

4. HARVEST Time of harvesting:

HS Mid-September HO Mid-October

NOTES: (1) SIL SCRF S-: Tubers treated with imazalil to guarantee no infection.

- (2) SIL SCRF SD: Tubers naturally infected with H. solari.
- (3) **BLK DOT** BD: Vermiculite, infected with *C. coccodes*, scattered on tubers at planting.

# Experimental diary:

07-Sep-93 : B : PK as (0:18:36) at 1390 kg.

29-Nov-93 : B : Ploughed.

08-Apr-94 : B : NPK as (12:20:20) at 1750 kg.

25-Apr-94 : B : Heavy spring-tine cultivated twice, rotary harrowed, ridged, potatoes hand planted. Ridges split back.

16-May-94 : B : Rotary ridged.

23-May-94 : B : Farmon PDQ at 2.0 l with Rotalin at 5.0 l in 200 l.
23-Jun-94 : B : Dithane 945 at 1.7 kg with Intracrop BLA at 0.2 l in 200 l.

## 94/R/P/4

### Experimental diary:

07-Jul-94: B: Dithane 945 at 1.7 kg with Pirimicarb 50 DG at 0.28 kg and Intracrop BLA at 0.2 l in 200 l.

25-Jul-94 : B : Ashlade Mancozeb FL at 2.5 1 with Intracrop BLA at 0.2 1 in 200 1.

15-Aug-94 : B : Ashlade Mancozeb FL at 2.5 1 with Intracrop BLA at 0.2 1 in 200 1.

30-Aug-94 : B : Ashlade Mancozeb FL at 2.5 1 with Intracrop BLA at 0.2 1 in 200 1.

24-Sep-94 : B : Haulm mechanically destroyed. 27-Sep-94 : T : HARVEST HS: Potatoes lifted. 25-Oct-94 : T : HARVEST HO: Potatoes lifted.

Previous crops: Lupins 1992, mustard and s. rape 1993.

NOTE: After harvest samples were assessed for silver scurf and black dot, further assessments were made after various storage regimes.

### TOTAL TUBERS TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

SIL SCRF	S-	SD	Mean
VARIETY			
ES	25.4	22.9	24.1
KE	56.5	51.5	54.0
Mean	41.0	37.2	39.1
BLK DOT	B-	BD	Mean
VARIETY			
ES	24.2	24.1	24.1
KE	56.3	51.8	54.0
Mean	40.2	37.9	39.1
BLK DOT	B-	BD	Mean
BLK DOT SIL SCRF	B-	BD	Mean
	B- 40.8	BD 41.1	Mean
SIL SCRF	-		
SIL SCRF	40.8	41.1	41.0
SIL SCRF S- SD Mean HARVEST	40.8	41.1 34.7	41.0 37.2
SIL SCRF S- SD Mean HARVEST VARIETY	40.8 39.7 40.2	41.1 34.7 37.9	41.0 37.2 39.1 Mean
SIL SCRF S- SD Mean HARVEST VARIETY ES	40.8 39.7 40.2 HS	41.1 34.7 37.9 HO	41.0 37.2 39.1 Mean 24.1
SIL SCRF S- SD Mean HARVEST VARIETY	40.8 39.7 40.2	41.1 34.7 37.9	41.0 37.2 39.1 Mean

94/R/P/4

# TOTAL TUBERS TONNES/HECTARE

\*\*\*\* Tables of means \*\*\*\*

HARV	EST	HS	HO	Mean		
SIL S	BCRF					
	S-	39.0	42.9	41.0		
	SD	35.6	38.8	37.2		
N	<b>l</b> ean	37.3	40.9	39.1		
HARV	/EST	HS	HO	Mean		
BLK	DOT					
	B-	37.9	42.5	40.2		
	BD	36.6	39.2	37.9		
N	Mean	37.3	40.9	39.1		
	SIL SCRF			SD		
VARIETY	BLK DOT	B-	BD	B-	BD	
ES		24.9		23.5	22.3	
KE		56.6	56.4	55.9	47.2	
	SIL SCRF			SD		
	HARVEST	HS	НО	HS	HO	
ES		24.1		22.5	23.2	
KE		53.9	59.2	48.6	54.4	
	BLK DOT	B-		BD		
	HARVEST	HS	НО	HS	НО	
ES		22.5	25.9		24.0	
KE		53.4	59.2	49.2	54.4	
	DIK DOM			22		
STI SCRE	BLK DOT HARVEST		110	BD	***	
S-	HARVEST	HS	HO	HS	НО	
SD SD		37.9				
50		37.9	41.4	33.2	36.2	
		BLK DOT	В-		22	
VARTETY	SIL SCRF		HS	по	BD	110
ES	S-	MARVEST	22.3	HO 27.5	HS 25.9	HO 25.8
25	SD		22.7		22.3	25.8
KE	S-		53.6	59.7	54.2	58.6
ALE.	SD		53.0	58.6	44.1	50.2
	50		55.4	50.0	44.1	50.2

## 94/R/P/4

# TOTAL TUBERS TONNES/HECTARE

\*\*\* Standard errors of differences of means \*\*\*

VARIETY	SIL SCRF	BLK DOT	HARVEST
0.90	0.90	0.90	0.90
VARIETY	VARIETY	SIL SCRF	VARIETY
SIL SCRF	BLK DOT	BLK DOT	HARVEST
1.28	1.28	1.28	1.28
SIL SCRF	BLK DOT	VARIETY	VARIETY
HARVEST	HARVEST	SIL SCRF	SIL SCRF
		BLK DOT	HARVEST
1.28	1.28	1.80	1.80
VARIETY	SIL SCRF	VARIETY	
BLK DOT	BLK DOT	SIL SCRF	
HARVEST	HARVEST	BLK DOT	
		HARVEST	
1.80	1.80	2.55	

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

 Stratum
 d.f.
 s.e.
 cv%

 BLOCK.WP
 30
 3.12
 8.0

PLOT AREA HARVESTED 0.00086