
Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1994



Full Table of Content

94/R/BN/7 Barnfield - Ley

Rothamsted Research

Rothamsted Research (1995) 94/R/BN/7 Barnfield - Ley; Yields Of The Field Experiments 1994, pp 32 - 34 - DOI: https://doi.org/10.23637/ERADOC-1-49

94/R/BN/7

BARNFIELD

Object: The experiment was designed to study the effects of organic and inorganic manures on continuous root crops. It was progressively modified to study effects on other crops.

Sections 1 and 2 the 11th year of grass/clover. The 20th year of grass on the rest of the experiment.

For previous years see 'Details' 1967 and 1973 and 74-93/R/BN/7.

Plot dimensions: 10.7 x 55.9.

Treatments to grass: All combinations of:-

Whole plots

1. MANURE Fertilizers and organic manures:

(D)	(D)
(D) PK	(D) P K
PKMG	P K (Na) Mg
P	P
PK	PK
PMG	P (Na) Mg
0	0

P: 35 kg P as triple superphosphate in 1974 and since 1987, single superphosphate in other years

K: 225 kg K as sulphate of potash

(Na): 90 kg Na as sodium chloride until 1973

Mg: 90 kg Mg as kieserite every fourth year since 1974 (sulphate of magnesia until 1973)

(D): Farmyard manure at 35 tonnes until 1975.

Quarter plots

2. N PERCUT	Nitrogen fertilizer in 1994 (kg N per cut) as 34.5% N,
	cumulative to previous dressings, and residues of
	forms of N previously each supplying 96 kg N per
	annum:

75	75, previously nitrate of soda, section 3
100	100, previously sulphate of ammonia, section 4
125	125, previously sulphate of ammonia + castor meal, section 5
150	150, previously castor meal, section 6

Castor meal last applied 1961, nitrate of soda and sulphate of ammonia until 1959.

Plus one plot MANURE KMG 100

94/R/BN/7

Treatments to grass/clover, sections 1 and 2 (not given nitrogen fertilizer):

MANURE Fertilizers and organic manures as for grass above, excluding KMG.

NOTES: (1) P, K and D treatments were applied to Sections 1 and 2 until 1980. None were applied subsequently until the resumption of P and K treatments only, from 1985.

- (2) Yields were not taken from section 2.
- (3) Only one cut was taken for yield as there was insufficient growth to justify a second cut.

Experimental diary:

19-Nov-93: T: Papplied.
23-Nov-93: T: Kapplied.
25-Nov-93: T: Mg applied.
18-Mar-94: T: Napplied to sections 3, 4, 5 and 6.
30-Mar-94: B: Flat rolled.
10-Jun-94: B: Cut.
11-Jun-94: B: Herbage cut on remainder of plot. Herbage removed.
14-Jun-94: T: Napplied to sections 3, 4, 5 and 6.
19-Oct-94: B: Topped. No yields taken.

NOTE: Herbage samples were taken for chemical analysis.

94/R/BN/7

GRASS

1ST CUT (10/6/94) DRY MATTER TONNES/HECTARE

**** Tables of means ****

N PERCUT	75	100	125	150	Mean
MANURE					
(D)	7.35	8.72	6.59	7.06	7.43
(D) PK	7.04	6.63	7.32	7.29	7.07
PKMG	6.66	7.64	7.61	7.68	7.40
P	3.85	3.84	3.95	5.27	4.23
PK	6.44	7.51	7.61	5.71	6.81
PMG	3.61	3.66	4.01	2.61	3.47
0	3.22	3.40	2.70	2.07	2.85
Mean	5.45	5.91	5.68	5.38	5.61

MANURE KMG 100 6.66

Grand mean 5.64

1ST CUT MEAN DM% 23.3

PLOT AREA HARVESTED 0.00568

GRASS/CLOVER

1ST CUT (10/6/94) DRY MATTER TONNES/HECTARE

***** Tables of means *****

MANURE (D) (D) PK PKMG P PK PMG 0 Mean 5.70 5.79 4.21 3.83 4.11 5.11 4.21 4.71

1ST CUT MEAN DM% 20.0

PLOT AREA HARVESTED 0.00155