Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1994



Full Table of Content

Tables of Means

Rothamsted Research

Rothamsted Research (1995) *Tables of Means ;* Yields Of The Field Experiments 1994, pp 7 - 7 - **DOI: https://doi.org/10.23637/ERADOC-1-49**

Harvest areas for cereals

On most of those cereal experiments which are harvested by combine the 'blank-row' technique is used to distinguish the areas taken for yield from the discard areas. For example when seed is drilled in 3 m wide plots in rows 12 cm apart appropriate coulters are prevented from sowing and 17 central rows are left for yield between pairs of blank rows. If the row-spacing is other than 12 cm a similar arrangement is used but with a different number of rows.

The ends of plots are separated from each other or from headlands by 1m fallow paths made after the crop has established.

The 'Area harvested' in the 'Yields', when the blank-row technique is used, is the product:-

number of rows harvested x distance between rows x length of rows.

If no rows are left blank and the plot is wider than the combine harvester so that discards are left uncut, the 'Area harvested' is the product:-

width of cutter bar x length of rows.

If the plot is narrower than the combine so that the whole area between paths is cut, the 'Area harvested' is the product:-

number of rows x distance between rows x length of rows.

We do not apply the adjustment used by some workers who take the harvested areas as width x length where each is measured to the centre of 'paths' up to a maximum of about 50 cm.

Tables of means

Tables of means are presented directly from computer output. Both factor and level names are presented in upper case characters. Vertical and horizontal lines are omitted e.g.:-

| FACTOR C | LEVEL C | L | LEVEL C | 2 | LEVEL C3 | |
|----------|---------|---------|-----------|------------|----------|----------|
| FACTOR B | LEVEL B | LEVEL B | 2 LEVEL B | 1 LEVEL B2 | LEVEL B1 | LEVEL B2 |
| FACTOR A | | | | | | |
| LEVEL A1 | * | * | * | * | * | * |
| LEVEL A2 | * | * | * | * | * | * |

Standard errors

NOTES: (1) This report gives standard errors of differences, not of means.

(2) Annotations (e.g. * min rep, max-min, max rep) to S.E.Ds are only explained the first time they occur in any experiment.