


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1994

[Full Table of Content](#)



94/W/CS/399 Cover Crops and N Cycling - W. Barley

Rothamsted Research

Rothamsted Research (1995) *94/W/CS/399 Cover Crops and N Cycling - W. Barley* ; Yields Of The Field Experiments 1994, pp 86 - 87 - DOI: <https://doi.org/10.23637/ERADOC-1-49>

94/R/CS/399

COVER CROPS AND N CYCLING

Object: To assess how effectively cover crops take up nitrogen and to assess how much of that nitrogen is subsequently made available to the following crop - Webbs.

Sponsors: P.R. Poulton, D.G. Christian, A.J. Macdonald.

The second year, w. barley.

Design: 3 blocks of 5 plots split into 2 sub-plots.

Whole plot dimensions: 9.0 x 12.0.

Treatments: All plots ploughed in autumn 1993 and sown to w. barley:

Whole plots

- | | |
|--------------------|---|
| 1. LAND TRT | Residues of cover crops, sown in autumn 1992, tumbledown and fallow. Plots ploughed in spring 1993 and sown to s. barley: |
| (FALLOWSB) | Fallow |
| (FO RA SB) | Forage rape |
| (RYE SB) | Rye |
| (TUMBDN SB) | Tumbledown |

Sub-plots

- | | |
|-------------|---|
| 2. N | Residues of nitrogen fertilizer (kg N) to s. barley 1993: |
| (0) | |
| (75) | |

plus one extra treatment

Whole plot

- | | |
|-----------------|--|
| 1. EXTRA | |
| (W BARLEY) | Residues of w. barley, sown in autumn 1992, taken to maturity: |

Sub-plot

- | | |
|-------------------|---|
| 2. N EXTRA | Residues of nitrogen fertilizer (kg N) to w. barley, sown in autumn 1992: |
| (0) | |
| (150) | |

94/R/CS/399

Experimental diary:

13-Sep-93 : B : Ploughed.
 08-Oct-93 : B : Rotary harrowed, Puffin, dressed Cerevax Extra, drilled at 350 seeds per m².
 12-Nov-93 : B : Optimol at 15 kg.
 19-Apr-94 : B : 34.5% N at 346 kg.
 19-Apr-94 : B : Briotril Plus 19/19 at 2.0 l with Hytane 500 SC at 2.0 l in 200 l.
 06-May-94 : B : Radar at 0.5 l with Standon Tridemorph 750 at 0.5 l in 200 l.
 31-May-94 : B : Mistral at 0.5 l with Radar at 0.5 l in 200 l.
 25-Jul-94 : B : Combine harvested.

NOTE: Soil water samples were taken for N analysis during the winter. Soil and crop samples were taken for N analysis during the summer.

GRAIN TONNES/HECTARE

***** Tables of means *****

	N	(0)	(75)	Mean
LAND TRT				
(FALLOWSB)		6.15	5.99	6.07
(FO RA SB)		6.37	6.01	6.19
(RYE SB)		6.74	6.87	6.80
(TUMBDN SB)		6.93	6.74	6.84
Mean		6.55	6.40	6.48

WINTER BARLEY

	N	(0)	(150)	Mean
N EXTRA		4.81	5.23	5.02

GRAND MEAN 6.19

*** Standard errors of differences of means ***

	LAND TRT	N	LAND TRT N & N EXTRA
	0.512	0.154	0.557
Except when comparing means with the same level(s) of			
CROPS			0.309

***** Stratum standard errors and coefficients of variation. *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	8	0.627	10.1
BLOCK.WP.SP	10	0.379	6.1
GRAIN MEAN DM%	88.3		
SUB PLOT AREA HARVESTED		0.00082	