


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1995

[Full Table of Content](#)



95/R/BK/1 Broadbalk - W. Wheat, Potatoes

Rothamsted Research

Rothamsted Research (1996) *95/R/BK/1 Broadbalk - W. Wheat, Potatoes* ; Yields Of The Field Experiments 1995, pp 11 - 17 - DOI: <https://doi.org/10.23637/ERADOC-1-50>

95/R/BK/1

BROADBALK

Object: To study the effects of organic and inorganic manures on continuous w. wheat. From 1968 two three-year rotations were included: potatoes, beans, w. wheat and fallow, w. wheat, w. wheat. In 1979 the first rotation was changed to fallow, potatoes, w. wheat. In 1980 the second rotation reverted to continuous w. wheat. Since 1985 part of the second rotation has been added to the first to extend the rotation to fallow, potatoes, w. wheat, w. wheat, w. wheat.

The 152nd year, w. wheat, fallow and potatoes.

For previous years see 'Details' 1967 and 1973, Station Report for 1966, pp. 229-231, Station Report for 1968, Part 2, Station Report for 1982, Part 2, pp. 5-44 and 74-94/R/BK/1.

Areas harvested:

Wheat:	Section	
	0	0.00351
	1	0.00645
	2,3,6 and 7	0.00533
	8 and 9	0.00561
Potatoes:	4	0.00348

Treatments:

Whole plots

PLOT

Fertilizers and organic manures:-

	Plot	Treatments until 1967	Treatments from 1968	Treatments from 1985
01DN4PK	01	-	D N2 P K	D N4 P K
21DN2	21	D	D N2	D N2
22D	22	D	D	D
030	03	None	None	None
05F	05	P K Na Mg	P K (Na) Mg	PK Mg
06N1F	06	N1 P K Na Mg	N1 P K (Na) Mg	N1 P K Mg
07N2F	07	N2 P K Na Mg	N2 P K (Na) Mg	N2 P K Mg
08N3F	08	N3 P K Na Mg	N3 P K (Na) Mg	N3 P K Mg
09N4F	09	N*1 P K Na Mg	N4 P K (Na) Mg	N4 P K Mg
10N2	10	N2	N2	N2
11N2P	11	N2 P	N2 P	N2 P
12N2PNA	12	N2 P Na	N2 P Na	N2 P Na
13N2PK	13	N2 P K	N2 P K	N2 P K
14N2PKMG	14	N2 P Mg	N2 P K Mg	N2 P K Mg
15N5F	15	N2 P K Na Mg	N3 P K (Na) Mg	N5 P K Mg
16N6F	16	N*2 P K Na Mg	N2 P K (Na) Mg	N6 P K Mg
17N0+3FH	17	N2 (A)	N2 1/2 (P K (Na) Mg)	N0+3 1/2 (PK Mg)+
18N1+3FH	18	P K Na Mg (A)	N2 1/2 (P K (Na) Mg)	N1+3 1/2 (PK Mg)+
19(C)	19	C	C	(C) (since 1989)
20N2KMG	20	N2 K Na Mg	N2 K (Na) Mg	N2 K Mg

(A) Alternating

95/R/BK/1

+ This change since 1980. Treatments shown are those to w. wheat; autumn N alternates. Potatoes receive N3 1/2 (PK Mg) on both Plots 17 and 18.

N1,N2,N3,N4,N5,N6: 48, 96, 144, 192, 240, 288 kg N (as sulphate of ammonia until 1967, except N* which was nitrate of soda. All as 'Nitro-Chalk' in spring from 1968 to 1985, as 34.5% N since 1986.)
 N0+3; N1+3: None in autumn + 144 kg N in spring; 48 kg N in autumn + 144 kg N in spring
 P: 35 kg P as triple superphosphate in 1974 and since 1988, single superphosphate in other years
 K: 90 kg K as sulphate of potash
 Na: 55 kg Na as sulphate of soda
 (Na): 16 kg Na as sulphate of soda until 1973
 Mg: 30 kg Mg annually to Plot 14, 35 kg Mg every third year to other plots since 1974. All as kieserite since 1974, previously as sulphate of magnesia annually
 D: Farmyard manure at 35 t
 (C): Castor meal to supply 96 kg N until 1988, none since
 F: P K (Na) Mg H: Half rate

Strips of sub-plots: Until 1967 wheat alone was grown on the experiment, with some bare fallowing. From 1968, ten strips of sub-plots (sections) were started with the following cropping:-

SECTION	1/W29	9/W37	0/W44	8/W1	6/W18	5/F	3/W3	7/W1	4/P	2/W2
Section	1	9	0*	8+	6**	5	3	7	4	2
Year										
1968	W	W	W	W	F	W	W	P	W	BE
1969	W	W	W	W	W	F	W	BE	P	W
1970	W	W	W	W	W	W	F	W	BE	P
1971	W	W	W	W	F	W	W	P	W	BE
1972	W	W	W	W	W	F	W	BE	P	W
1973	W	W	W	W	W	W	F	W	BE	P
1974	W	W	W	W	F	W	W	P	W	BE
1975	W	W	W	W	W	F	W	BE	P	W
1976	W	W	W	W	W	W	F	W	BE	P
1977	W	W	W	W	F	W	W	P	W	BE
1978	W	W	W	W	W	F	W	BE	P	W
1979	W	W	W	W	W	W	F	W	P	F
1980	W	W	W	W	W	W	W	F	W	P
1981	W	W	W	F	W	W	W	P	F	W
1982	W	W	W	W	W	W	W	W	P	F
1983	W	W	W	W	W	W	W	F	W	P
1984	W	W	W	W	W	W	W	P	F	W
1985	W	W	W	W	W	F	W	W	P	W
1986	W	W	W	W	W	P	F	W	W	W
1987	W	W	W	W	W	W	P	W	W	F
1988	W	W	W	F	W	W	W	F	W	P
1989	W	W	W	W	W	W	W	P	F	W
1990	W	W	W	W	W	F	W	W	P	W

95/R/BK/1

SECTION	1/W29	9/W37	0/W44	8/W1	6/W18	5/F	3/W3	7/W1	4/P	2/W2
Section	1	9	0*	8+	6**	5	3	7	4	2
Year										
1991	W	W	W	W	W	P	F	W	W	W
1992	W	W	W	W	W	W	P	W	W	F
1993	W	W	W	W	W	W	W	F	W	P
1994	W	W	W	F	W	W	W	P	F	W
1995	W	W	W	W	W	F	W	W	P	W

W = w. wheat, P = potatoes, BE = s. beans, F = fallow

* Straw incorporated since autumn 1986. ** No sprays except weedkillers since 1985. + No weedkillers.

NOTES: (1) For a fuller record of treatments see 'Details' etc.
 (2) From autumn 1975 to autumn 1986, chalk was applied at 2.9 t each autumn to all plots in sets of Sections on a three-year cycle. Year 1: Sections 1,2,3. Year 2: Sections 6,7,8,9. Year 3: Sections 0,4,5. From autumn 1988 until autumn 1992 a five-year cycle was used. Year 1: Sections 1,3. Year 2: Sections 2,8. Year 3: Sections 7,9. Year 4: Sections 4,6. Year 5: Sections 0,5. None applied since autumn 1991.

Experimental diary:

All sections:

- 10-Oct-94 : T : Mg and Na applied.
- 11-Oct-94 : T : P and K applied.
- 13-Oct-94 : T : Farmyard manure applied.
- : B : Ploughed and furrow pressed.

Cropped sections:

W. wheat:

- 24-Aug-94 : T : Straw chopped (section 0 only).
- 30-Aug-94 : T : Straw baled (sections 1, 2, 3, 5, 6, and 9).
- 12-Sep-94 : T : Weeds topped (section 8 only).
- 12-Oct-94 : T : Autumn N treatments applied.
- 16-Nov-94 : T : Rotary harrowed, Apollo, dressed New Kotol, drilled at 380 seeds per m².
- 21-Mar-95 : T : Stefes IPU at 3.0 l with Stomp 400 at 3.3 l in 200 l (except section 8).
- 11-Apr-95 : T : Spring N treatments applied.
- 28-Apr-95 : T : Halo at 2.0 l with Mallard 750 EC at 0.5 l and New 5C Cycocel at 2.8 l in 200 l (except section 6).
- 19-Jun-95 : T : Sprint at 1.5 l in 300 l (except section 6).
- 05-Jul-95 : T : Pirimicarb 50 DG at 280 g in 200 l (except section 6).
- 18-Jul-95 : T : Pulled wild oats by hand.
- 27-Jul-95 : T : Roundup at 6.0 l with Mixture B at 2.94 l in 150 l (except section 8).
- 06-Aug-95 : T : Combine harvested.

Potatoes:

- 12-Sep-94 : T : Weeds topped.
- 11-Apr-95 : T : N treatments applied.
- 24-Apr-95 : T : Heavy spring-tine cultivated twice, rotary harrowed, planted Estima, undressed.
- 04-May-95 : T : Rotary ridged.

95/R/BK/1

Experimental diary:

Potatoes:

- 01-Jun-95 : T : Basagran at 3.0 l in 400 l.
- 20-Jun-95 : T : Ashlade Maneb Flowable at 2.75 l in 200 l.
- 30-Jun-95 : T : Ashlade Maneb Flowable at 2.75 l in 200 l.
- 19-Jul-95 : T : Ashlade Maneb Flowable at 2.75 l in 300 l.
- 01-Aug-95 : T : Ashlade Maneb Flowable at 2.75 l in 300 l.
- 14-Aug-95 : T : Super-Tin 4L at 560 ml with Intracrop BLA at 200 ml in
200 l.
- 24-Aug-95 : T : Haulm pulverised.
- 15-Sep-95 : T : Lifted.

Fallow:

- 30-Aug-94 : T : Straw baled.
- 24-Apr-95 : T : Heavy spring-tine cultivated twice.
- 09-Jun-95 : T : Cultivated by rotary-grubber.
- 17-Jul-95 : T : Spring-tine cultivated.

NOTE: Samples of grain and straw from sections 1 and 7 and samples of potato tubers were taken for chemical analysis.

95/R/BK/1 W. WHEAT

GRAIN TONNES/HECTARE

***** Tables of means *****

SECTION PLOT	7/W1	8/W1	2/W2	3/W3	6/W18	1/W29	9/W37	0/W44
01DN4PK	9.65	*	9.08	8.88	8.17	*	*	*
21DN2	9.62	7.91	8.27	7.19	7.04	8.44	7.38	6.55
22D	7.43	6.95	6.78	5.91	5.66	7.36	5.27	6.09
030	1.91	2.32	0.78	0.61	1.82	1.13	1.10	1.54
05F	1.56	4.86	0.57	1.04	1.70	1.32	0.86	1.48
06N1F	4.57	4.82	3.03	3.13	3.73	3.60	4.33	3.79
07N2F	6.71	5.69	4.81	3.30	4.99	5.25	5.32	4.67
08N3F	7.75	5.93	6.31	4.14	5.76	6.08	5.56	5.88
09N4F	8.02	5.71	6.30	5.19	6.07	6.16	6.67	5.86
10N2	4.87	4.54	4.04	2.45	2.99	2.56	2.67	2.29
11N2P	5.20	3.22	4.07	3.24	2.70	3.11	2.74	2.52
12N2PNA	5.27	3.64	3.99	3.60	3.83	3.27	3.04	3.64
13N2PK	6.19	5.42	4.56	3.87	4.71	4.73	4.86	4.54
14N2PKMG	5.77	6.32	4.58	3.79	5.23	5.35	4.58	5.38
15N5F	7.75	5.58	7.37	6.58	6.75	6.94	6.48	7.21
16N6F	8.13	6.18	7.48	7.11	7.14	7.37	7.58	7.56
17N0+3FN	7.51	6.07	5.95	5.16	6.29	6.34	6.22	5.46
18N1+3FN	7.80	5.85	7.15	6.03	6.95	7.03	6.84	6.45
19 (C)	1.63	4.83	0.82	1.64	1.75	2.40	1.49	1.70
20NKMG	*	*	*	*	*	3.25	*	3.10

GRAIN MEAN DM% 89.1

95/R/BK/1 W. WHEAT

STRAW TONNES/HECTARE

***** Tables of means *****

SECTION	7/W1	1/W29
PLOT		
01DN4PK	5.61	*
21DN2	4.16	4.83
22D	2.10	3.32
030	0.29	0.37
05F	0.08	0.31
06N1F	0.90	0.92
07N2F	1.81	1.93
08N3F	2.50	2.59
09N4F	2.58	2.72
10N2	1.40	1.57
11N2P	1.48	1.56
12N2PNA	1.11	1.33
13N2PK	1.24	1.64
14N2PKMG	1.03	1.54
15N5F	2.54	2.85
16N6F	3.37	2.86
17N0+3FN	1.91	1.94
18N1+3FN	2.56	2.33
19 (C)	0.14	0.27
20NKMG	*	1.32

STRAW MEAN DM% 93.2

95/R/BK/1 POTATOES

***** Tables of means *****

PLOT	TOTAL TUBERS	% WARE
	TONNES/ HECTARE	3.81 CM (1.5 INCH) RIDDLE
01DN4PK	8.9	76.2
21DN2	11.2	70.4
22D	10.1	68.9
030	1.8	24.6
05F	3.5	30.7
06N1F	6.7	43.4
07N2F	6.7	47.9
08N3F	10.2	67.1
09N4F	10.1	74.1
10N2	2.4	20.7
11N2P	3.5	14.9
12N2PNA	4.3	22.9
13N2PK	7.7	51.5
14N2PKMG	10.9	69.5
15N5F	11.4	74.5
16N6F	11.7	74.4
17N3FH	6.3	51.7
18N3FH	9.3	64.1
19 (C)	3.7	22.4