


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1995

[Full Table of Content](#)



95/R/LN/1 Weed Types in Linseed - Linseed

Rothamsted Research

Rothamsted Research (1996) *95/R/LN/1 Weed Types in Linseed - Linseed* ; Yields Of The Field Experiments 1995, pp 156 - 157 - DOI: <https://doi.org/10.23637/ERADOC-1-50>

95/R/LN/1

LINSEED

WEED TYPES IN LINSEED

Object: To study the effects of three weed species on the growth and yield of linseed - Hoosfield Old 4-course.

Sponsor: P.J.W. Lutman.

Design: 3 blocks of 15 plots.

Plot dimensions: 3.0 x 10.0.

Treatments:

SPECDEN	Weed species and density (average number of established plants per m ²):
0	None (triplicated)
C1	Cultivated oats (<i>Avena sativa</i> cv. Dula), 10.7
C2	Cultivated oats, 30.7
C3	Cultivated oats, 86.1
C4	Cultivated oats, 174.0
K1	Knotgrass (<i>Polygonum aviculare</i>), 20.3
K2	Knotgrass, 47.2
K3	Knotgrass, 74.4
K4	Knotgrass, 145.0
F1	Fat hen (<i>Chenopodium album</i>), 0
F2	Fat hen, 0
F3	Fat hen, 0
F4	Fat hen, 23.3

- NOTES:** (1) Target weed sowing rates and densities were as follows:
Cultivated oats (seeds per m²): C1 20, C2 60, C3 180 and C4 360.
Knotgrass (plants per m²): K1 20, K2 80, K3 180 and K4 320.
Fat hen (plants per m²): F1 40, F2 80, F3 160 and F4 320.
- (2) Weeds on plots with treatments F1, F2, F3 failed to establish. These plots have been omitted from the analysis.

Experimental diary:

- 06-Sep-94 : B : PK as (0:20:32) at 1317 kg.
07-Sep-94 : B : Dolomite at 5.0 t.
16-Dec-94 : B : Ploughed.
19-Apr-95 : B : Heavy spring-tine cultivated. Rotary harrowed.
20-Apr-95 : **T** : **SPECDEN** C1, C2, C3, C4: Dula, dressed Rappor, drilled.
 : **T** : **SPECDEN** K1, K2, K3, K4: Knotgrass broadcast.
 : **T** : **SPECDEN** F1, F2, F3, F4: Fat hen broadcast.
 : B : Rotary harrowed, Antares, dressed Prelude 20 LF, drilled
 at 700 seeds per m².

95/R/LN/1

Experimental diary:

21-Apr-95 : B : Rolled.
04-May-95 : B : Part irrigated, 25 mm.
05-May-95 : B : Part irrigated, 25 mm.
10-May-95 : B : Decis at 200 ml in 300 l.
11-May-95 : B : 34.5% N at 232 kg.
15-Aug-95 : B : Standon Diquat at 3.0 l with Vassgro Spreader at 400 ml
in 400 l.
22-Aug-95 : B : Hand harvested.

Previous crops: W. oats 1993, w. wheat 1994.

NOTE: Crop and weed densities were assessed in May. Crop and weeds were sampled in June and August to measure dry matter.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

SPECDEN	
0	1.11
C1	0.91
C2	0.59
C3	0.29
C4	0.18
K1	0.89
K2	0.77
K3	0.61
K4	0.58
F4	0.98
Mean	0.76

*** Standard errors of differences of means ***

SPECDEN	
0.125	min.rep
0.102	max-min

SPECDEN
max-min 0 v any of the remainder
min.rep Any of the remainder

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	24	0.154	20.1
GRAIN MEAN DM%	not measured		
PLOT AREA HARVESTED	0.00020		