


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1995

[Full Table of Content](#)



95/W/P/2 Double Cropping and Nematodes - Potatoes

Rothamsted Research

Rothamsted Research (1996) *95/W/P/2 Double Cropping and Nematodes - Potatoes* ; Yields Of The Field Experiments 1995, pp 166 - 168 - DOI: <https://doi.org/10.23637/ERADOC-1-50>

95/W/P/2

POTATOES

DOUBLE CROPPING AND NEMATODES

Object: To assess the population dynamics of *Globodera rostochiensis* in double cropped potatoes - Woburn, Lansome III.

Sponsors: K. Evans, P.D. Halford.

Design: 4 randomised blocks of 6 plots.

Whole plot dimensions: 3.0 x 12.0.

Treatments:

1. **NEMACIDE[1]** Nematicide applied to first double crop:

- None
OX Oxamyl

2. **NEMACIDE[2]** Nematicide applied to second double crop:

- None
OX Oxamyl

plus 2 extra plots

3. **NEMMAIN** Nematicide applied to main crop:

- None
OX Oxamyl

Experimental diary:

08-Sep-94 : B : PK as (0:20:32) at 781 kg
21-Sep-94 : B : Ploughed
21-Mar-95 : **T** : **NEMACIDE[1]** OX: Vydate at 40 kg.
 : **T** : Double crop: 34.5% N at 522 kg. PK as (0:24:24) at
 1000 kg. Rotary cultivated, Maris Bard, dressed
 Rizolex, planted.
23-Mar-95 : B : Campbell's Linuron 45% Flowable at 1.9 l with Gramoxone
 at 4.0 l in 400 l.
24-Mar-95 : **T** : Double crop: Covered with polythene.
01-May-95 : **T** : **NEMMAIN** OX: Vydate at 40 kg.
 : **T** : Main crop: 34.5% N at 522 kg. PK as (0:24:24) at
 1000 kg. Rotary cultivated, Estima, dressed Rizolex,
 planted.
26-May-95 : **T** : Double crop: Polythene removed .
08-Jun-95 : **T** : Double crop: Gramoxone 100 at 4.0 l in 200 l sprayed
 between rows.
20-Jun-95 : **T** : Double crop: Hand weeded.
05-Jul-95 : B : Ashlade Maneb Flowable at 2.5 l in 300 l.
11-Jul-95 : B : Irrigated 12.5 mm.
12-Jul-95 : B : Irrigated 12.5 mm.
17-Jul-95 : **T** : Double crop: Potatoes lifted.

95/W/P/2

Experimental diary:

- 18-Jul-95 : T : **NEMACIDE[2]** OX: Vydate at 40 kg.
: T : Double crop: PK as (0:24:24) at 1000 kg, 27% N at 667 kg. Rotary cultivated.
- 19-Jul-95 : T : Double crop: Rotary cultivated, Estima, dressed Rizolex, planted.
- 21-Jul-95 : B : Irrigated 20 mm.
- 24-Jul-95 : B : Ashlade Maneb Flowable at 2.5 l in 400 l.
- 25-Jul-95 : B : Irrigated 12.5 mm.
- 11-Aug-95 : B : Irrigated 12.5 mm.
- 12-Aug-95 : B : Ashlade Maneb Flowable at 2.5 l in 400 l.
- 14-Aug-95 : B : Irrigated 12.5 mm.
- 17-Aug-95 : B : Irrigated 12.5 mm.
- 23-Aug-95 : T : Main crop: Potatoes lifted.
: T : Double crop: Harvest at 4.0 l in 300 l sprayed between rows.
- 31-Aug-95 : T : Double crop: Irrigated 12.5 mm.
- 13-Sep-95 : T : Double crop: Super-Tin 4L at 560 ml in 300 l.
- 11-Oct-95 : T : Double crop: Super-Tin 4L at 560 ml in 300 l.
- 25-Oct-95 : T : Double crop: Potatoes lifted.

Previous crops: Potatoes 1993, S. barley 1994.

NOTE: Soil samples were taken at intervals through the growing season for nematode counts. Plant samples of roots and tops were taken to measure vegetative production and nematode infestation.

95/W/P/2

TOTAL TUBERS TONNES/HECTARE

***** Tables of means *****

NEMACIDE [2]	-	OX	Mean
NEMACIDE [1]			
-	14.0	20.1	17.0
OX	21.9	23.8	22.9
Mean	18.0	21.9	20.0
NEMMAIN			
-	14.0		
OX	24.1		
Mean	19.1		

GRAND MEAN 19.7

*** Standard errors of differences of means ***

NEMACIDE [1]	NEMACIDE [2]	NEMACIDE [1] NEMACIDE [2] & NEMMAIN
1.89	1.89	2.67

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	15	3.78	19.2
PLOT AREA HARVESTED	0.00150		