


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1995

[Full Table of Content](#)



### 95/W/CS/347 Green Crops for Set-aside - W. Wheat , S. Wheat

#### Rothamsted Research

Rothamsted Research (1996) 95/W/CS/347 *Green Crops for Set-aside - W. Wheat , S. Wheat ; Yields Of The Field Experiments 1995*, pp 76 - 78 - DOI: <https://doi.org/10.23637/ERADOC-1-50>

95/W/CS/347

**GREEN CROPS FOR SET-ASIDE**

**Object:** To obtain information on the establishment and maintenance of sown crops and unsown vegetation in three-year and five-year set-aside. Effects on soil nitrate and leaching after ploughing are also studied - Woburn, Horsepool Lane Close II.

**Sponsors:** R.D. Prew, E.T.G. Bacon, D.P. Yeoman, M.V. Hewitt, J.F. Jenkyn, R.J. Gutteridge.

**Design:** 3 randomised blocks of 6 plots split into 2 x 2 criss-cross.

**Whole plot dimensions:** 6.5 x 26.0.

The sixth year, w. and s. sown wheat.

For previous years see 90-94/W/CS/347.

**Treatments:**

Test phase (first test crop after 5-year treatment phase)

Whole plots

1. **PREVCROP** Previous crops, cumulative 1990 to 1994:
  - (RY LF) Ryegrass, cuttings left in situ
  - (RY+CL LF) Ryegrass + clover, cuttings left in situ
  - (RY+CL RE) Ryegrass + clover, cuttings removed
  - (RY+N RE) Ryegrass given 100 kg N in spring, cuttings removed
  - (TU LF) Tumbledown, natural regrowth, cuttings left in situ
  - (ARABLE) Arable sequence w. wheat, w. wheat, w. oats, w. wheat, w. oats

Sub-plots (**WHEAT** split-plots, **N** criss-cross)

2. **WHEAT** Time of ploughing and drilling:
  - W Winter
  - S Spring
3. **N** Fertilizer nitrogen applied in spring:
  - NO None
  - N OPT Optimum

**NOTES:** (1) N OPT was given 40 kg N early (W in March, S in June) and the following rates later (W in April, S in July)

	(RY LF)	(RY+CL LF)	(RY+CL RE)	(RY+N RE)	(TU LF)	(ARABLE)
W	155	135	135	160	140	170
S	120	120	120	120	130	155

(2) The three blocks after 3-year set-aside were discontinued after the second test crop in 1994.

95/W/CS/347

**Experimental diary:**

05-Oct-94 : T : WHEAT W: Ploughed.  
07-Oct-94 : T : WHEAT W: Rolled. Rotary harrowed, Cadenza, dressed Rappor, drilled at 300 seeds per m<sup>2</sup>.  
01-Dec-94 : B : Stomp 400 at 3.3 l with Stefes IPU at 3.0 l in 200 l.  
16-Mar-95 : T : WHEAT W: N N OPT: 27% N at 148 kg.  
22-Mar-95 : T : WHEAT S: Ploughed.  
30-Mar-95 : T : WHEAT S: Rotary harrowed.  
31-Mar-95 : T : WHEAT S: Rotary harrowed, Cadenza, dressed Rappor, drilled at 500 seeds per m<sup>2</sup>.  
28-Apr-95 : T : WHEAT W, N N OPT: Nitrogen treatments applied as 27% N.  
          : T : WHEAT S: Rotary harrowed, Cadenza, dressed Cerevax, re-drilled at 500 seeds per m<sup>2</sup>.  
02-May-95 : T : WHEAT W: Halo at 2.0 l in 200 l.  
25-May-95 : T : WHEAT S: Cadenza, dressed Rappor, re-drilled at 900 seeds per m<sup>2</sup>.  
26-May-95 : T : PREVCROP (RY+CL RE), (RY+N RE): Corrective P and K applied.  
01-Jun-95 : T : WHEAT W: Cyclone at 1.0 l in 300 l.  
15-Jun-95 : T : WHEAT S: Ally at 30 g in 300 l.  
16-Jun-95 : T : WHEAT S, N N OPT: 27% N at 148 kg.  
29-Jun-95 : B : Pirimicarb 50 DG at 280 g in 300 l.  
19-Jul-95 : T : WHEAT S, N N OPT: Nitrogen treatments applied as 27% N.  
05-Aug-95 : T : WHEAT W: Combine harvested.  
22-Sep-95 : T : WHEAT S: Combine harvested.

- NOTES:** (1) The spring wheat was re-drilled twice following the failure of the first and second sowings, and yields were negligible and are not presented.
- (2) Soil mineral nitrogen was measured in autumn and spring. Weeds were counted in November and July. Numbers of ears of wheat were estimated in July.

95/W/CS/347 PLOTS 19-36

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

	N	NO	N OPT	Mean
<b>PREVCROP</b>				
(RY LF)		1.70	6.74	4.22
(RY+CL LF)		4.03	8.29	6.16
(RY+CL RE)		3.72	7.17	5.45
(RY+N RE)		2.15	7.12	4.63
(TU LF)		1.44	7.64	4.54
(ARABLE)		1.72	7.24	4.48
Mean		2.46	7.37	4.91

\*\*\* Standard errors of differences of means \*\*\*

	PREVCROP	N	PREVCROP	N
	0.707	0.304	0.884	
Except when comparing means with the same level(s) of				
<b>PREVCROP</b>			0.752	
<b>N</b>			0.886	

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP1.WP2	10	0.923	18.8

GRAIN MEAN DM% not measured

SUB-PLOT AREA HARVESTED 0.00235 or 0.00323