


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1996

[Full Table of Content](#)



---

### 96/W/CS/347 Green Crops for Set-aside - W. Wheat

#### Rothamsted Research

Rothamsted Research (1997) *96/W/CS/347 Green Crops for Set-aside - W. Wheat* ; Yields Of The Field Experiments 1996, pp 75 - 77 - DOI: <https://doi.org/10.23637/ERADOC-1-51>

96/W/CS/347

### GREEN CROPS FOR SET-ASIDE

**Object:** To obtain information on the establishment and maintenance of sown crops and unsown vegetation in three-year and five-year set-aside. Effects on soil nitrate and leaching after ploughing are also studied - Woburn, Horsepool Lane Close II.

**Sponsors:** E.T.G. Bacon, D.P. Yeoman, M.V. Hewitt, J.F. Jenkyn, R.J. Gutteridge.

**Design:** 3 randomised blocks of 6 plots split into 2 x 2 criss-cross.

**Whole plot dimensions:** 6.5 x 26.0.

The seventh year, w. wheat.

For previous years see 90-95/W/CS/347.

**Treatments:**

Test phase (second test crop after 5-year treatment phase)

Whole plots

- |             |   |
|-------------|---|
| 1. PREVCROP | Previous crop, cumulative 1990 to 1994:                       |
| (RY LF)     | Ryegrass, cuttings left in situ                               |
| (RY+CL LF)  | Ryegrass + clover, cuttings left in situ                      |
| (RY+CL RE)  | Ryegrass + clover, cuttings removed                           |
| (RY+N RE)   | Ryegrass given 100 kg N in spring, cuttings removed           |
| (TU LF)     | Tumbledown, natural regrowth, cuttings left in situ           |
| (ARABLE)    | Arable sequence w. wheat, w. wheat, w. oats, w.wheat, w. oats |

Sub-plots (**WHEAT** split-plots, **N** criss-cross)

- |          |  |
|----------|--|
| 2. WHEAT | Time of ploughing and drilling previous w. wheat crop:                         |
| (W)      | Winter 1994  |
| (S)      | Spring 1995  |
| 3. N     | Fertilizer nitrogen (kg N) applied in spring cumulative to previous dressings: |
| NO       | None   |
| N OPT    | Optimum, 40 early and 160 later  |

**Experimental diary:**

- 03-Oct-95 : B : Roundup at 4.0 l in 300 l.  
12-Oct-95 : B : Ploughed.  
19-Oct-95 : B : Rotary harrowed, Cadenza, dressed Beret 050FS, drilled at 375 seeds per m<sup>2</sup>.  
22-Nov-95 : B : Stomp 400 SC at 2.5 l with Dagger at 1.5 l in 200 l.  
11-Mar-96 : T : N N OPT: 34.5% N at 116 kg.

96/W/CS/347

**Experimental diary:**

30-Apr-96 : B : Halo at 1.5 l in 200 l.  
 02-May-96 : T : N N OPT: 27.5% N at 581 kg.  
 06-Jun-96 : B : Halo at 2.0 l with Mallard 750 EC at 0.5 l in 300 l.  
 20-Aug-96 : B : Combine harvested.

**NOTE:** Soil nitrogen was measured in autumn. Weeds were counted and estimates of ground cover were made in spring.

**GRAIN TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

	N	NO	NOPT	Mean
<b>PREVCROP</b>				
(RY LF)		5.14	8.48	6.81
(RY+CL LF)		6.24	9.44	7.84
(RY+CL RE)		6.45	8.58	7.52
(RY+N RE)		4.80	8.56	6.68
(TU LF)		4.36	8.11	6.23
(ARABLE)		4.53	8.58	6.56
Mean		5.25	8.63	6.94
<b>WHEAT</b>	(W)	(S)		Mean
<b>PREVCROP</b>				
(RY LF)		6.41	7.21	6.81
(RY+CL LF)		7.48	8.20	7.84
(RY+CL RE)		7.02	8.02	7.52
(RY+N RE)		6.51	6.85	6.68
(TU LF)		6.29	6.18	6.23
(ARABLE)		6.20	6.91	6.56
Mean		6.65	7.23	6.94
<b>WHEAT</b>	(W)	(S)		Mean
<b>N</b>				
NO		4.76	5.75	5.25
NOPT		8.55	8.71	8.63
Mean		6.65	7.23	6.94
<b>PREVCROP</b>	<b>WHEAT</b>	(W)	(S)	
	<b>N</b>	NO	NOPT	NO
(RY LF)		4.60	8.23	5.69
(PC)		5.56	9.40	6.93
(PCR)		5.76	8.27	7.14
(RY+N RE)		4.56	8.46	5.04
(TU LF)		4.07	8.51	4.65
(ARABLE)		3.99	8.41	5.06
				NOPT
				8.74
				9.48
				8.90
				8.66
				7.71
				8.75

96/W/CS/347

GRAIN TONNES/HECTARE

\*\*\* Standard errors of differences of means \*\*\*

	PREVCROP	WHEAT	PREVCROP WHEAT
	1.015	0.166	1.054
Except when comparing means with the same level(s) of PREVCROP			0.406

	PREVCROP* N	WHEAT* N	PREVCROP* WHEAT N
	1.218	0.646	1.260
Except when comparing means with the same level(s) of PREVCROP	0.740		0.844
WHEAT		0.631	
N	1.060	0.187	1.108
PREVCROP.WHEAT			0.769
PREVCROP.N			0.457
WHEAT.N			1.108

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP1	10	1.243	17.9
BLOCK.WP1.SP	12	0.497	7.2
BLOCK.WP1.WP2	10	0.532	7.7
BLOCK.WP1.SP.WP2	12	0.366	5.3

GRAIN MEAN DM% 88.1

SUB-PLOT AREA HARVESTED 0.00236 or 0.00325