Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1997



Full Table of Content

97/R/CS/483 Severe Take-all in Wheat - S. Wheat

Rothamsted Research

Rothamsted Research (1998) 97/R/CS/483 Severe Take-all in Wheat - S. Wheat; Yields Of The Field Experiments 1997, pp 100 - 101 - DOI: https://doi.org/10.23637/ERADOC-1-53

97/R/CS/483

SEVERE TAKE-ALL IN WHEAT

Object: To create severe take-all (Gaeumannomyces graminis) in winter wheat by applying inoculum artificially to a preceding spring wheat - Summerdells I.

Sponsors: G.L. Bateman, R.J. Gutteridge.

Design: 4 randomised blocks of 14 plots.

Plot dimensions: 3.0 x 10.0.

Treatments: -

INOCULTN Inoculum:

None

TA Take-all inoculum on sterile oat grain

NOTE: Each treatment appears seven times in each block.

Experimental diary:

08-Jul-96 : B : Deep tine cultivated with vibrating tines 60 cm apart and 45 cm deep.

10-Jul-96 : B : Rolled.

25-Oct-96 : B : Ploughed.

11-Mar-97 : B : Rotary harrowed, spring-tine cultivated twice.

: T : INOCULTN TA: Inoculated oat grain broadcast.

12-Mar-97: B: Rotary harrowed, Chablis, undressed, drilled at 400 seeds per m^2 .

14-Mar-97 : B : Rolled.

12-May-97 : B : 34.5% N at 370 kg.

22-May-97 : B : Campbell's CMPP at 2.1 1 with Vindex at 1.0 1 in 300 1.

05-Sep-97 : B : Combine harvested.

Previous crops: W. and S. rape 1995, set-aside 1996.

NOTE: Take-all patches were assessed in July.

97/R/CS/483

GRAIN TONNES/HECTARE

***** Tables of means *****

INOCULTN

7.29 6.26 TA

6.77 Mean

*** Standard errors of differences of means ***

INOCULTN

0.142

***** Stratum standard errors and coefficients of variation *****

Stratum

d.f.

s.e.

CV%

BLOCK.WP

51

0.529 7.8

GRAIN MEAN DM% 84.5

PLOT AREA HARVESTED 0.00240