


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1997

[Full Table of Content](#)



97/W/WW/1 Sulphur, Variety and Nitrogen - W. Wheat

Rothamsted Research

Rothamsted Research (1998) *97/W/WW/1 Sulphur, Variety and Nitrogen - W. Wheat* ; Yields Of The Field Experiments 1997, pp 104 - 105 - DOI: <https://doi.org/10.23637/ERADOC-1-53>

97/W/WW/1

WINTER WHEAT

SULPHUR, VARIETY AND QUALITY

Object: To measure yield and quality response to sulphur fertilizer on three varieties of wheat - Woburn, Butt Close.

Sponsors: S.P. McGrath, F. Zhao.

Design: 3 randomised blocks of 3 x 2 x 3 plots

Plot dimensions: 3.0 x 12.0.

Treatments: All combinations of:-

1. VARIETY

H	Hereward dressed Sibutol
S	Spark dressed Sibutol
R	Rialto dressed Panoctine

2. NITROGEN Nitrogen fertilizer (kg N) as 27.5% N:

N1	180
N2	230

3. SULPHUR Sulphur fertilizer (kg S) as gypsum (17.5% S):

S-	None
S1	20
S2	100

Experimental diary:

01-Oct-96 : B : Ploughed.
02-Oct-96 : B : Rolled. Rotary harrowed.
03-Oct-96 : T : **VARIETY** H, S, R: Varieties drilled at 350 seeds per m².
06-Dec-96 : B : Panther at 2.0 l in 200 l.
20-Mar-97 : T : **SULPHUR** S1, S2: Gypsum applied at 114 and 571 kg respectively.
20-Mar-97 : B : 27.5% N at 145 kg.
03-Apr-97 : T : **NITROGEN** N1, N2: 27.5% N applied at 509 and 691 kg respectively.
23-May-97 : B : Standon Fluroxypyr at 0.75 l with Halo at 2.0 l in 300 l.
08-Aug-97 : B : Barclay Gallup at 2.0 l in 300 l.
14-Aug-97 : B : Combine harvested.

Previous crops: Potatoes 1995, s. barley 1996.

NOTE: Plant samples were taken in May and June for measurement of sulphur and nitrogen content. Harvest samples of straw and grain were also analysed for sulphur and nitrogen. Grain samples from selected plots were tested for bread making quality.

97/W/WW/1

GRAIN TONNES/HECTARE

***** Tables of means *****

VARIETY NITROGEN	H	S	R	Mean
N1	3.30	5.15	3.89	4.12
N2	3.94	4.09	4.29	4.11
Mean	3.62	4.62	4.09	4.11

SULPHUR NITROGEN	S-	S1	S2	Mean
N1	4.21	3.99	4.15	4.12
N2	3.27	4.29	4.77	4.11
Mean	3.74	4.14	4.46	4.11

SULPHUR VARIETY	S-	S1	S2	Mean
H	3.34	3.66	3.86	3.62
S	4.20	5.18	4.49	4.62
R	3.67	3.58	5.03	4.09
Mean	3.74	4.14	4.46	4.11

NITROGEN	SULPHUR VARIETY	S-	S1	S2
N1	H	3.46	3.17	3.28
	S	5.24	5.03	5.18
	R	3.92	3.76	3.99
N2	H	3.23	4.14	4.44
	S	3.16	5.32	3.79
	R	3.41	3.40	6.07

*** Standard errors of differences of means ***

VARIETY	NITROGEN	SULPHUR	VARIETY NITROGEN
0.383	0.313	0.383	0.542

VARIETY SULPHUR	NITROGEN SULPHUR	VARIETY NITROGEN SULPHUR
0.664	0.542	0.939

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	34	1.150	28.0
GRAIN MEAN DM%	87.9	PLOT AREA HARVESTED	0.00220