

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1997

[Full Table of Content](#)



97/W/RAW/1 Diagnosis of S Deficiency - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1998) *97/W/RAW/1 Diagnosis of S Deficiency - W. Oilseed Rape* ; Yields Of The Field Experiments 1997, pp 130 - 131 - DOI: <https://doi.org/10.23637/ERADOC-1-53>

97/W/RAW/1

WINTER OILSEED RAPE
DIAGNOSIS OF S DEFICIENCY

Object: To study the effects of rates of sulphur on the yield and sulphur content of winter oilseed rape - Woburn, Far Field II.

Sponsors: F. Zhao, S.P. McGrath.

Design: 4 randomised blocks of 6 plots.

Whole plot dimensions: 3.0 x 12.0.

Treatments:

SULPHUR	Sulphur as gypsum (17.5% S) kg S:
S0	0
S1	5
S2	10
S3	20
S4	40
S5	80

Experimental diary:

- 02-Sep-96 : T : SULPHUR 5, 10, 20, 40, 80: Gypsum applied at 28.6, 57.1, 114, 229, 457 kg respectively.
- 02-Sep-96 : B : Drilled Apex, dressed Lindex-Plus FS at 6.6 kg per ha.
- 24-Jul-97 : B : Reglone at 3.0 l with Vassgro Non-ionic at 400 ml in 400 l.
- 05-Aug-97 : B : Combine harvested.

Previous crops: Potatoes 1995, w. wheat 1996.

NOTE: Plant samples were taken on four occasions between March and May to measure total sulphur content, sulphate-S and glutathione.

97/W/RAW/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

SULPHUR

S0	4.51
S1	4.17
S2	4.39
S3	4.44
S4	4.27
S5	4.33

Mean 4.35

*** Standard errors of differences of means ***

SULPHUR

0.151

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
---------	------	------	-----

BLOCK.WP	15	0.213	4.9
----------	----	-------	-----

GRAIN MEAN DM% 90.6

PLOT AREA HARVESTED 0.00288