

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1997

[Full Table of Content](#)



97/R/LP/10 Genotype, Row Spacing and Seed Rate - Lupins

Rothamsted Research

Rothamsted Research (1998) *97/R/LP/10 Genotype, Row Spacing and Seed Rate - Lupins* ; Yields Of The Field Experiments 1997, pp 146 - 148 - DOI: <https://doi.org/10.23637/ERADOC-1-53>

97/R/LP/10

LUPINS

GENOTYPE, ROW SPACING AND SEED RATE

Object: To test seed rate and row spacing on the structure and performance of existing determinate and new dwarf-determinate genotypes - Stackyard.

Sponsors: I. Shield, G.F.J. Milford, J.E. Leach.

Design: 3 randomised blocks of 4 x 2 x 2 plots.

Whole plot dimensions: 9.0 x 9.0.

Treatments: All combinations of :-

1. GENOTYPE

70	CH304/70
73	CH304/73
12	DTN 12
20	DTN 20

2. ROW SPAC Row spacing, cm:

R1	11
R2	36

3. SEED RAT Seed rate, seeds per m²:

S1	40
S2	80

Experimental diary:

25-Jun-96 : B : Ploughed and furrow pressed.
26-Jul-96 : B : Rolled.
12-Sep-96 : B : Spring-tine cultivated.
13-Sep-96 : B : Rotary harrowed.
 : T : Genotypes undressed drilled at 40 and 80 seeds per m² respectively.
23-Sep-96 : B : Stomp 400 SC at 5.0 l in 294 l. Spannit at 1.5 l in 294 l.
04-Oct-96 : B : Irrigated 25 mm.
08-Nov-96 : B : Carbetamex at 3.0 kg with MSS Simazine 50 FL at 2.3 l.
 Decis at 300 ml in 200 l.
12-Dec-96 : B : Rovral Flo at 1.0 l in 200 l, Standon Tebuconazole at 0.5 l in 200 l.
01-May-97 : B : Compass at 3.0 l in 200 l.
18-Jun-97 : B : Mistral at 1.0 l in 300 l.
08-Jul-97 : B : Danadim Dimethoate 40 at 850 ml in 300 l.
05-Sep-97 : B : Harvest at 3.0 l in 400 l.
23-Sep-97 : B : Combine harvested.

Previous crops: W. wheat 1995, set-aside 1996.

97/R/LP/10

- NOTES: (1) Plant populations were assessed in autumn, January, April and at harvest. Leaf and branch numbers were assessed in June, time of flowering was noted, light interception was measured frequently during the growing season. Dry matter and nitrogen accumulation was assessed at intervals on certain plots. After harvest oil and nitrogen content and grain density was measured.
- (2) Most plots of **GENOTYPE** 73 failed and it has been omitted from the analysis.

GRAIN TONNES/HECTARE

***** Tables of means *****

ROW SPAC	R1	R2	Mean
GENOTYPE			
70	2.21	2.04	2.12
12	2.35	1.74	2.05
20	2.22	1.74	1.98
Mean	2.26	1.84	2.05
SEED RAT	S1	S2	Mean
GENOTYPE			
70	1.52	2.73	2.12
12	1.44	2.66	2.05
20	2.03	1.93	1.98
Mean	1.66	2.44	2.05
SEED RAT	S1	S2	Mean
ROW SPAC			
R1	1.88	2.64	2.26
R2	1.45	2.24	1.84
Mean	1.66	2.44	2.05
GENOTYPE	SEED RAT	S1	S2
	ROW SPAC		
70	R1	1.56	2.86
	R2	1.48	2.59
12	R1	1.58	3.12
	R2	1.30	2.19
20	R1	2.50	1.94
	R2	1.56	1.93

97/R/LP/10

GRAIN TONNES/HECTARE

*** Standard errors of differences of means ***

GENOTYPE	ROW SPAC	SEED RAT	GENOTYPE
			ROW SPAC
0.218	0.178	0.178	0.309
GENOTYPE	ROW SPAC	GENOTYPE	
SEED RAT	SEED RAT	ROW SPAC	
		SEED RAT	
0.309	0.252	0.437	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	22	0.535	26.1
GRAIN MEAN DM%	85.9		
PLOT AREA HARVESTED	0.00216		