


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1997

[Full Table of Content](#)



97/R/CS/302 Eyespot Resistance to Mbc - W. Wheat

Rothamsted Research

Rothamsted Research (1998) *97/R/CS/302 Eyespot Resistance to Mbc - W. Wheat* ; Yields Of The Field Experiments 1997, pp 59 - 60 - DOI: <https://doi.org/10.23637/ERADOC-1-53>

97/R/CS/302

EYESPOT RESISTANCE TO MBC

Object: To study the development of resistance to MBC fungicides in eyespot and the ability of resistant strains to survive, spread and infect - Meadow.

Sponsor: G.L. Bateman.

The 13th year, w. wheat.

For previous years see 85-93,95-96/R/CS/302

Design: 2 randomised blocks of 4 plots split into 6 sub-plots.

Whole plot dimensions: 12.0 X 24.0.

Sub-plot dimensions: 4.5 x 6.0.

Treatments: All combinations of:-

Whole plots

1. FUNGICIDE	Fungicide applied cumulatively 1985-93 and 1995-97:
NONE	None
CARB	Carbendazim at 0.25 kg
PRO	Prochloraz at 0.40 kg (0.50 kg in 1993, 1995-1997)
CARB+PRO	Carbendazim and prochloraz as above

Sub-plots

2. EYE INOC	Eyespot inoculum, applied in first year only:
NATURAL	Natural background population (duplicated)
W 19R 1S	Inoculated with wheat strains in proportion 19 resistant to one sensitive
W 1R 19S	As above but one resistant to 19 sensitive
R 19R 1S	Inoculated with rye strains, 19 resistant to one sensitive
R 1R 19S	As above but one resistant to 19 sensitive

NOTE: The inoculum was colonized on oat seed and broadcast in October, 1984.

Experimental diary:

19-Sep-96 : B : Ploughed and furrow pressed.
03-Oct-96 : B : Rotary harrowed, Hereward, dressed Beret Gold, drilled at 380 seeds per m².
05-Dec-96 : B : Isoproturon 500 at 2.6 l with Stomp 400 SC at 3.1 l in 200 l.
11-Mar-97 : B : 34.5% N at 118 kg.
20-Mar-97 : T : **FUNGICIDE** CARB+PRO: Barclay Eyetak at 1.1 l with Tripart Defensor FL at 0.5 l in 200 l.
 : T : **FUNGICIDE** PRO: Barclay Eyetak at 1.1 l in 200 l.

97/R/CS/302

Experimental diary:

20-Mar-97 : T : FUNGCIDE CARB: Tripart Defensor FL at 0.5 l in 200 l.
 04-Apr-97 : T : 34.5% N at 463 kg.
 16-Apr-97 : T : FUNGCIDE CARB+PRO: Barclay Eyetak at 1.1 l with Tripart
 Defensor FL at 0.5 l in 200 l.
 : T : FUNGCIDE PRO: Barclay Eyetak at 1.1 l in 200 l.
 : T : FUNGCIDE CARB: Tripart Defensor FL at 0.5 l in 200 l.
 18-Apr-97 : B : Starane 2 at 0.75 l with Barclay Holdup at 2.3 l in 300 l.
 18-Aug-97 : B : Combine harvested.

NOTE: Plant samples were taken in July from **EYE INOC NATURAL** plots for assessment of stem base diseases.

GRAIN TONNES/HECTARE

***** Tables of means *****

EYE INOC FUNGCIDE	NATURAL	W 19R 1S	W 1R 19S	R 19R 1S	R 1R 19S	Mean
NONE	4.94	5.77	3.64	4.56	5.75	4.93
CARB	4.78	5.13	5.01	5.34	5.16	5.04
PRO	5.20	4.27	5.41	5.46	5.46	5.17
CARB+PRO	5.62	5.28	4.98	4.95	6.13	5.43
Mean	5.13	5.11	4.76	5.08	5.63	5.14

*** Standard errors of differences of means ***

EYE INOC	FUNGCIDE*	EYE INOC
0.299	0.598	min.rep
0.259	0.518	max-min

EYE INOC
 max-min NATURAL v any of the remainder
 min.rep Any of the remainder

* Within the same level of **FUNGCIDE** only

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP.SP	24	0.598	11.6
GRAIN MEAN DM%	88.1		
SUB-PLOT AREA HARVESTED	0.00144		