

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1997

[Full Table of Content](#)



---

### 97/R/CS/476 Fungicide Sequences and Take-all - W. Wheat

#### Rothamsted Research

Rothamsted Research (1998) *97/R/CS/476 Fungicide Sequences and Take-all - W. Wheat* ; Yields Of The Field Experiments 1997, pp 94 - 95 - DOI: <https://doi.org/10.23637/ERADOC-1-53>

97/R/CS/476

### FUNGICIDE SEQUENCES AND TAKE-ALL

**Object:** To determine the effects of fungicidal seed treatments on take-all (*Gaeumannomyces graminis*) development in w.wheat - Long Hoos IV 4.

**Sponsors:** G.L. Bateman, J.F. Jenkyn.

The first year, w. wheat.

**Design:** 4 randomised blocks of 2 x 2 x 2.

**Plot dimensions:** 3.0 x 10.0.

**Treatments:** All combinations of:-

1. **FUNG97**                      Fungicidal seed dressing to the 1997 crop:  
    F97                          Seed dressed  
    -97                          None
2. **FUNG98**                      Fungicidal seed dressing to the 1998 crop:  
    F98                          Seed dressed  
    -98                          None
3. **FUNG99**                      Fungicidal seed dressing to the 1999 crop:  
    F99                          Seed dressed  
    -99                          None

**NOTE:** The seed dressing is under commercial development, composition undisclosed.

**Experimental diary:**

- 17-Oct-96 : B : Spring-tine cultivated, rotary harrowed, Brigadier, dressed as treatment, drilled at 380 seeds per m<sup>2</sup>.
- 28-Jan-97 : B : Panther at 2.0 l in 200 l.
- 11-Mar-97 : B : 34.5% N at 118 kg.
- 14-Apr-97 : B : 34.5% N at 463 kg.
- 16-Apr-97 : B : Deloxil at 1.5 l with Dow Shield at 0.35 l in 200 l.
- 30-May-97 : B : Folicur at 0.5 l in 300 l.
- 19-Aug-97 : B : Combine harvested.

Previous crops: S. wheat 1995, linseed 1996.

**NOTE:** Plant samples were taken in April to assess take-all on the roots and in July to assess root and stem base diseases. Soil samples were taken after harvest and used in bioassays to measure take-all infectivity.

97/R/CS/476

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

FUNG97

F97 10.31

-97 10.02

Mean 10.17

\*\*\* Standard errors of differences of means \*\*\*

FUNG97

0.225

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	27	0.637	6.3
GRAIN MEAN DM%	90.0		
PLOT AREA HARVESTED	0.00240		