

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1998

[Full Table of Content](#)



98/R/CS/483 Severe Take-all in Wheat - W. Wheat

Rothamsted Research

Rothamsted Research (1999) *98/R/CS/483 Severe Take-all in Wheat - W. Wheat* ; Yields Of The Field Experiments 1998, pp 108 - 109 - DOI: <https://doi.org/10.23637/ERADOC-1-52>

98/R/CS/483

SEVERE TAKE-ALL IN WHEAT

Object: To create severe take-all (*Gaeumannomyces graminis*) in winter wheat by applying inoculum artificially to spring wheat and then test seed dressings on w. wheat - Summerdells I.

Sponsors: G.L. Bateman, R.J. Gutteridge.

The second year, w. wheat.

For previous year see 97/R/CS/483

Design: 4 randomised blocks of 2 x 7 plots.

Plot dimensions: 3.0 x 10.0.

Treatments:-

All combinations of:

1. **INOCULTN** Inoculum applied in 1997:
 - None
 - TA Take-all inoculum

2. **SEED TRT** Seed dressing in 1998:
 - None
 - R1 CR 21529 at 200 ml per 100 kg
 - R2 CR 21529 at 300 ml per 100 kg
 - R3 CR 21529 at 450 ml per 100 kg
 - NR1 CR 21529 (Non-P) at 200 ml per 100 kg
 - NR3 CR 21529 (Non-P) at 450 ml per 100 kg
 - B Baytan Flowable at 200 ml per 100 kg

NOTE: Treatments CR 21529 and CR 21529 (Non-P) are under commercial development, composition disclosed in confidence.

Experimental diary:

- 10-Sep-97 : B : PK as (0:20:32) at 1250 kg.
- 01-Oct-97 : B : Heavy spring-tine cultivated twice.
- 01-Oct-97 : T : Rotary harrowed, Hereward, dressed as treatment, drilled at 400 seeds per m².
- 03-Oct-97 : B : Rolled.
- 16-Oct-97 : B : Draza at 5.5 kg.
- 13-Jan-98 : B : Unite A at 0.125 l with Unite B at 1.0 l and LI-700 at 1.0 l in 200 l.
- 17-Feb-98 : B : 34.5% N at 120 kg.
- 29-Apr-98 : B : 34.5% N at 460 kg.
- 18-Aug-98 : B : Combine harvested.

NOTE: Plant populations, growth and take-all were assessed in March. Take-all and stem base diseases were assessed in June.

98/R/CS/483

GRAIN TONNES/HECTARE

***** Tables of means *****

INOCULTN SEED TRT	-	TA	Mean
-	5.78	5.82	5.80
R1	6.03	6.25	6.14
R2	6.14	6.58	6.36
R3	5.59	6.67	6.13
NR1	6.75	5.68	6.21
NR3	6.22	6.70	6.46
B	5.68	6.29	5.99
Mean	6.03	6.28	6.16

*** Standard errors of differences of means ***

INOCULTN	SEED TRT	INOCULTN SEED TRT
0.227	0.425	0.601

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	39	0.850	13.8
GRAIN MEAN DM%	88.0		
PLOT AREA HARVESTED	0.00228		