


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1998

[Full Table of Content](#)



---

### 98/W/CS/491 Efficiency of S Fertilisers - W. Wheat

#### Rothamsted Research

Rothamsted Research (1999) *98/W/CS/491 Efficiency of S Fertilisers - W. Wheat* ; Yields Of The Field Experiments 1998, pp 112 - 113 - DOI: <https://doi.org/10.23637/ERADOC-1-52>

98/W/CS/491

### EFFICIENCY OF S FERTILIZERS

**Object:** To measure the effect of different forms of sulphur on yields of wheat and a subsequent rape crop - Woburn, Stackyard III.

**Sponsors:** F.J. Zhao, S.P. McGrath.

The first year, w. wheat.

**Design:** 4 randomised blocks of 4 x 2 + 1.

**Plot dimensions:** 4.0 x 12.0.

**Treatments:-**

1. S FORM	Form of sulphur:
T+A	Tiger 90 and ammonium sulphate
AS	Ammonium sulphate
T90	Tiger 90
NAS	Sodium thiosulphate

2. APP TIME	Time of application:
SB	To the seedbed
MAR	17-Mar-98

**EXTRA**

-	None
---	------

**NOTE:** Nitrogen fertilizer was applied to balance the nitrogen supplied by ammonium sulphate, this was 26 kg to the seedbed and to give a total of 180 kg N in March.

**Experimental diary:**

29-Sep-97 : B : Ploughed.  
01-Oct-97 : B : Rolled.  
02-Oct-97 : B : Rotary harrowed.  
03-Oct-97 : T : **APP TIME** SB: Sulphur treatments applied to give 30 kg S.  
                  : T : Balancing nitrogen applied as 27% N.  
03-Oct-97 : B : Riband, dressed Sibutol, drilled at 385 seeds per m<sup>2</sup>.  
13-Nov-97 : B : Stomp 400 SC at 2.0 l with Isoproturon 500 at 1.0 l and  
                  Cyperkill 10 at 0.25 in 200 l.  
17-Mar-98 : T : **APP TIME** MAR: Sulphur treatments applied to give 30 kg S.  
                  : T : Balancing nitrogen applied as 27% N.  
04-May-98 : B : Alto 100 SL at 0.6 l in 200 l. Tripart Brevis at 1.5 l  
                  in 200 l.  
31-May-98 : B : Opus at 0.8 l in 200 l.  
12-Jun-98 : B : Folicur at 0.3 l with Bavistin DF at 0.3 kg in 200 l.  
13-Aug-98 : B : Combine harvested.

Previous crops: W. rye 1996, potatoes 1997.

98/W/CS/491

**NOTE:** Soils were sampled in October and August and analysed for sulphur.  
Plants were sampled in April, July and August for sulphur and nitrogen.

# GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

APP TIME	SB	MAR	Mean
<b>S FORM</b>			
T+A	9.41	8.98	9.19
AS	9.37	9.38	9.37
T90	9.11	9.15	9.13
NAS	9.19	9.30	9.24
Mean	9.27	9.20	9.24
<b>EXTRA</b>	9.37		

GRAND MEAN 9.25

\*\*\* Standard errors of differences of means \*\*\*

S FORM	APP TIME	S FORM APP TIME & EXTRA
0.177	0.125	0.250

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	24	0.353	3.8

GRAIN MEAN DM% 87.9

PLOT AREA HARVESTED 0.00230