
Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1998



Full Table of Content

# 98/R/RN/22 Crop Rotation - W. Oats

### **Rothamsted Research**

Rothamsted Research (1999) 98/R/RN/22 Crop Rotation - W. Oats; Yields Of The Field Experiments 1998, pp 51 - 51 - DOI: https://doi.org/10.23637/ERADOC-1-52

#### 98/R/RN/22

#### CROP ROTATIONS

Object: To test combinable break crops and their effect on following wheat crops. New crop species and winter sown varients of established species will be tested to determine optimal break crop selection for rotations that maximise first wheat yields and minimise inputs - Great Field I/II.

Sponsor: I.F. Shield, M.V. Hewitt, R.W. Payne.

The first year, w.oats.

Design: 2 blocks of 42 plots split into 4 sub-plots.

Whole plot dimensions:  $4.5 \times 10.0$ .

#### Treatments:

None (preparatory year)

#### Experimental diary:

```
15-Sep-97 : B : Deep tine cultivated with vibrating tines 60 cm apart and 45 cm deep.

18-Sep-97 : B : Chalk at 4.0 t.

19-Sep-97 : B : Ploughed.

13-Oct-97 : B : Rotary harrowed, Image, dressed Anchor, drilled at 350 seeds per m².

13-Feb-98 : B : 34.5% N at 116 kg.

01-Apr-98 : B : Mistral at 0.75 l in 200 l.

27-Apr-98 : B : 34.5% N at 155 kg.

27-Apr-98 : B : Rovral Flo at 3.0 l in 200 l.

12-May-98 : B : Tripart Brevis at 2.25 l with Headland Enhance LF at
```

40 ml in 200 l.

25-Jun-98 : B : Corbel at 1.0 1 in 200 1.

06-Aug-98: T: Combine harvested.

NOTE: The whole site was sown to oats in the preparatory year. Yields were taken for each sub-plot and a yield map produced. Lodging occured in

## GRAIN TONNES/HECTARE

many plots.

Grand mean 6.30

SUB-PLOT AREA HARVESTED 0.00230