


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1998

[Full Table of Content](#)



### 98/R/CS/472 Cereals and Seed Treatments - W. Wheat, W. Barley

#### Rothamsted Research

Rothamsted Research (1999) *98/R/CS/472 Cereals and Seed Treatments - W. Wheat, W. Barley ; Yields Of The Field Experiments 1998*, pp 93 - 94 - DOI: <https://doi.org/10.23637/ERADOC-1-52>

98/R/CS/472

# CEREALS AND SEED TREATMENTS

**Object:** To test seed treatment fungicides on root and stem base diseases of winter wheat and barley - Highfield IV/Road Piece East.

**Sponsors:** W.A.J.M. Dawson, G.L. Bateman, J.F. Jenkyn.

The second year, w. wheat and w. barley.

**Design:** 4 randomised blocks of 8 x 2.

**Plot dimensions:** 3.0 x 10.0.

**Treatments:** All combinations of:-

1. FUNGICIDE	Seed dressing:	
	1997	1998
(-)-	None	None
(E)-	CR21528	None
(-)E	None	CR21529
(E)E	CR21528	CR21528
(B)-	CR21529	None
(-)B	None	CR21529
(B)B	CR21529	CR21529

## 2. CROP

WW	Winter wheat
BW	Winter barley

**NOTE:** Fungicides CR21528 and CR21529 are under commercial development, composition disclosed in confidence.

## Experimental diary:

17-Sep-97 : B : Scythe LC at 3.0 l with Vassgro Non Ionic at 100 ml in 200 l.  
 18-Sep-97 : B : Topped.  
 24-Sep-97 : B : Ploughed and furrow pressed.  
 26-Sep-97 : T : CROP BW: Rotary harrowed, Pipkin, dressed as treatment, drilled at 350 seeds per m<sup>2</sup>.  
                   : T : CROP WW: Rotary harrowed, Brigadier, dressed as treatment, drilled at 380 seeds per m<sup>2</sup>.  
 06-Jan-98 : B : Atlas Fieldgard at 2.6 l with Stomp 400 SC at 3.3 l in 200 l.  
 13-Feb-98 : B : 34.5% N at 116 kg. Grasp at 1.4 l with Isoguard at 2.0 l and Output at 0.75 l in 200 l.  
 25-Feb-98 : B : Manganese sulphate at 3.0 kg with Tern 750 EC at 0.75 l in 200 l.  
 31-Mar-98 : T : CROP BW: Campbell's Carbendazim 50% Flowable at 0.5 l in 220 l.  
 28-Apr-98 : B : 34.5% N at 400 kg.  
 08-May-98 : B : Ally at 20 g with Starane 2 at 0.5 l in 200 l.  
 14-May-98 : B : Carbate Flowable at 0.5 l with Opus at 1.0 l in 200 l.

98/R/CS/472

**Experimental diary:**

05-Jun-98 : B : Corbel at 1.0 l in 200 l.  
 20-Jul-98 : B : CROP BW: Combine harvested.  
 11-Aug-98 : B : CROP WW: Combine harvested.

**NOTE:** Plant samples were taken in January, March and June to assess root and stem base diseases.

**GRAIN TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

CROP FUNGICIDE	WW	BW	Mean
(-) -	7.59	7.40	7.50
(E) -	7.99	7.51	7.75
(-) E	9.14	7.33	8.23
(E) E	8.47	7.85	8.16
(B) -	8.00	7.90	7.95
(-) B	9.32	7.26	8.29
(B) B	8.74	7.43	8.08
Mean	8.36	7.51	7.93

\*\*\* Standard errors of differences of means \*\*\*

FUNGICIDE	CROP	FUNGICIDE CROP
0.296		0.419 min.rep
0.257	0.148	0.363 max-min
		0.296 max.rep

**FUNGICIDE**

max.rep - only  
 max-min - v any of the remainder  
 min.rep Any of the remainder

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	47	0.593	7.5
GRAIN MEAN DM%	88.5		
PLOT AREA HARVESTED	0.00228		