

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1999

[Full Table of Content](#)



99/R/WW/4 Fusarium Study - W. Wheat

Rothamsted Research

Rothamsted Research (2000) *99/R/WW/4 Fusarium Study - W. Wheat* ; Yields Of The Field Experiments 1999, pp 120 - 121 - DOI: <https://doi.org/10.23637/ERADOC-1-34>

99/R/WW/3

Experimental diary:

19-May-99 : B : : tm) Sypex at 1.5 l in 200 l.
 : B : : tm) Enhance at 80 ml in 200 l.
21-May-99 : B : : Irrigated, 11 mm.
28-May-99 : T : : Nitrogen applied as 34.5% N (GS 51).
 : B : : tm) Opus at 1.0 l in 200 l.
 : B : : tm) Sipcam UK Rover 500 at 1.0 l in 200 l.
16-Jun-99 : B : : Folicur at 0.25 l in 100 l.
06-Aug-99 : B : : Combine harvested.

Previous crops: W. rape and turnip rape 1997, w. oats 1998.

GRAIN TONNES/HECTARE

***** Tables of means *****

NITROGEN

-	4.13
A	7.70
B	9.12
C	10.36
D	9.45
E	10.65
F	11.27
G	10.23
H	11.05
I	11.24
J	10.67
K	5.95
L	7.60
M	8.72
N	6.80
O	8.39
P	10.46
Q	11.03

Mean 9.16

*** Standard errors of differences of means ***

NITROGEN

0.248

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	34	0.304	3.3
GRAIN MEAN DM%	86.0		
PLOT AREA HARVESTED	0.00352		

99/R/WW/4

WINTER WHEAT

FUSARIUM STUDY

Object: To assess effects of different inocula and inoculum sources on ear blight development in mist-irrigated w. wheat - Little Knott I.

Sponsors: G.L. Bateman, J.F. Jenkyn, R.J. Gutteridge.

Design: 3 randomised blocks of 6 x 2 plots

Whole plot dimensions: 6.0 x 8.5.

Treatments: All combinations of:-

1. **T** Seed and *Fusarium* inocula:
 - D O Diseased seed, no inoculum
 - H O Healthy seed, no inoculum
 - H A Healthy seed, inoculated with *Fusarium culmorum* to soil in spring
 - H B Healthy seed, inoculated with *Fusarium culmorum* to ears at anthesis
 - H C Healthy seed, inoculated with *Microdochium nivale* to ears at anthesis

2. **IRRIGATN** Irrigation:
 - None
 - I Irrigated

Experimental diary:

- 14-Sep-98 : B : : Ploughed and furrow pressed.
- 19-Sep-98 : B : : Rotary harrowed.
- 23-Sep-98 : T : D : Combination drilled, Charger, homesaved, at 380 seeds/m² with the Accord drill.
- : T : H : Combination drilled, Charger, tr. Beret Gold, at 380 seeds/m² with the Accord drill.
- 23-Feb-99 : B : : tm)Isoguard at 1.0 l in 200 l.
- : B : : tm)Unite A at 0.125 l in 200 l.
- : B : : tm)Unite B at 1.0 l in 200 l.
- : B : : tm)Cropoil at 1.0 l in 200 l.
- 12-Mar-99 : B : : 34.5% N at 145 kg.
- 18-Mar-99 : B : : Triple superphosphate at 300 kg.
- 31-Mar-99 : T : H A : Inoculated with colonised sterile oat grain, at 1.1 kg to the central 6 x 3 metre area of each plot.
- 15-Apr-99 : B : : 34.5% N at 435 kg.
- 31-May-99 : T : I : Plots misted during anthesis, four at a time, for 2 days, between 31-May and 8-Jun, immediately after spore inoculum applied to ears.
- : T : H B : Inoculated between 31-May and 6-Jun, by spraying spore suspensions onto plots during anthesis.
- : T : H C : Inoculated 31-May to 6-Jun, by spraying spore suspensions onto plots during anthesis.
- 30-Jul-99 : B : : Combine harvested.