

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1999

[Full Table of Content](#)



99/R/CS/302 Eyespot Resistance to Mbc - W. Wheat

Rothamsted Research

Rothamsted Research (2000) *99/R/CS/302 Eyespot Resistance to Mbc - W. Wheat* ; Yields Of The Field Experiments 1999, pp 52 - 53 - DOI: <https://doi.org/10.23637/ERADOC-1-34>

99/R/RN/22

Experimental diary:

W. lupins:

16-Oct-98 : T : tm) Stomp 400 SC at 5.0 l in 220 l.
 : T : tm) Decis at 300 ml in 220 l.
08-Feb-99 : T : Carbetamex at 3.0 kg in 220 l.
17-Mar-99 : T : tm) Folicur at 0.5 l in 200 l.
 : T : tm) Rovral Flo at 1.0 l in 200 l.

NOTE: Yields are presented by crop not by **ROTATION**. Winter peas and lupins failed and therefore there are no yields.

***** Tables of means *****

WINTER CROPS	GRAIN TONNES/ HECTARE	GRAIN MEAN DM%
OATS	7.44	89.4
WHEAT	10.30	89.0
RAPE	3.96	90.2
LINSEED	1.40	93.1
BEANS	5.39	78.7

SUB-PLOT AREA HARVESTED 0.00230

99/R/CS/302

EYESPOT RESISTANCE TO MBC

Object: To study the development of resistance to MBC fungicides in eyespot and the ability of resistant strains to survive, spread and infect - Meadow.

Sponsor: G.L. Bateman.

The 15th year, w. wheat.

For previous years see 85-93,95-98/R/CS/302

Design: 2 randomised blocks of 4 plots split into 6 sub-plots.

Whole plot dimensions: 12.0 x 24.0.

Sub-plot dimensions: 4.5 x 6.0.

Treatments: All combinations of:-

Whole plots

- | | |
|---------------------|---|
| 1. FUNGICIDE | Fungicide applied cumulatively 1985-93 and 1995-99: |
| NONE | None |
| CARB | Carbendazim at 0.25 kg |
| PRO | Prochloraz at 0.40 kg (0.50 kg in 1993, 1995-1999) |
| CARB+PRO | Carbendazim and prochloraz as above |

Sub-plots

- | | |
|--------------------|---|
| 2. EYE INOC | Eyespot inoculum, applied in first year only: |
| NATURAL | Natural background population (duplicated) |
| W 19R 1S | Inoculated with wheat strains in proportion 19 resistant to one sensitive |
| W 1R 19S | As above but one resistant to 19 sensitive |
| R 19R 1S | Inoculated with rye strains, 19 resistant to one sensitive |
| R 1R 19S | As above but one resistant to 19 sensitive |

NOTE: The inoculum was colonized on oat seed and broadcast in October, 1984.

Experimental diary:

18-Sep-98	: B :	: Ploughed and furrow pressed, started.
19-Sep-98	: B :	: Ploughed and furrow pressed, completed.
21-Sep-98	: B :	: Combination drilled, Hereward, tr. Beret Gold, at 380 seeds/m ² with the Accord drill.
01-Dec-98	: B :	: tm)Amazon at 1.0 l in 200 l.
	: B :	: tm)Isogard at 2.0 l in 200 l.
	: B :	: tm)Stefes Cypermethrin 2 at 250 ml in 200 l.
	: B :	: tm)Cropoil at 1.0 l in 200 l.
04-Mar-99	: B :	: 34.5% N at 145 kg.
17-Mar-99	: T : CARB	: Stefes C-Flo 2 at 0.5 l in 200 l.
	: T : CARB+PRO	: Sportak 45 EW at 1.1 l with Stefes C-Flo 2 at 0.5 l in 200 l.