


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1999

[Full Table of Content](#)



99/R/CS/326 and 99/W/CS/326 Amounts of Straw - W. Wheat

Rothamsted Research

Rothamsted Research (2000) *99/R/CS/326 and 99/W/CS/326 Amounts of Straw - W. Wheat* ; Yields Of The Field Experiments 1999, pp 62 - 65 - DOI: <https://doi.org/10.23637/ERADOC-1-34>

99/R/CS/323

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNG99 GROUP	NONE	FLUQUIN	Mean
I	9.50	9.97	9.73
II	9.05	9.69	9.37
III	9.00	9.81	9.41
Mean	9.18	9.82	9.50

*** Standard errors of differences of means ***

GROUP	FUNG99	GROUP FUNG99
0.160	0.131	0.227

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	64	0.555	5.8

GRAIN MEAN DM% 87.6

PLOT AREA HARVESTED 0.00227

99/R/CS/326 and 99/W/CS/326

AMOUNTS OF STRAW

Object: To study the effects of different amounts of straw, incorporated into the soil, on w. wheat - Rothamsted (R) Great Knott III, Woburn (W) Far Field I.

Sponsors: M.J. Glendining, P.C. Brookes.

The 13th year, w. wheat.

For previous years see 87-98/R & W/CS/326.

Design: 4 randomised blocks of 4 plots (R).
3 randomised blocks of 4 plots (W).

Whole plot dimensions: 3.0 x 13.5 (R).
3.0 x 14.5 (W).

Treatments:

STRAW Amounts of straw incorporated into the seedbed (t per ha 85% DM), cumulative to previous annual dressings:

		R	W
NONE	None	-	-
NORMAL	Normal	5.8	5.6
2 NORMAL	Twice normal	11.6	11.2
4 NORMAL	Four times normal	23.2	22.4

Experimental diary:

Great Knott III (R):

20-Aug-98 : T : NONE : Straw removed.
: T : NORMAL, 2 NORMAL, 4 NORMAL : Straw applied and chopped.
08-Oct-98 : B : : Ploughed.
13-Oct-98 : B : : Rotary harrowed. Drilled Hereward, tr. Sibutol, at
380 seeds/m² with the Nordsten drill.
: B : : Hardy at 10.0 kg.
15-Oct-98 : B : : Hardy at 7.5 kg.
04-Nov-98 : B : : Decoy at 7.5 kg.
10-Mar-99 : B : : tm)Isoguard at 1.0 l in 200 l.
: B : : tm)Unite A at 0.125 l in 200 l.
: B : : tm)Unite B at 1.0 l in 200 l.
: B : : tm)Cropoil at 1.0 l in 200 l.
15-Mar-99 : B : : 34.5% N at 145 kg.
12-Apr-99 : B : : 34.5% N at 435 kg.
01-May-99 : B : : Opus at 0.7 l in 200 l.
29-May-99 : B : : tm)Folicur at 1.0 l in 100 l.
: B : : tm)Sipcam UK Rover 500 at 1.0 l in 100 l.
17-Jun-99 : B : : Folicur at 0.25 l in 100 l.
23-Aug-99 : B : : Combine harvested.

99/R/CS/326 and 99/W/CS/326

Experimental diary:

Far Field I (W):

17-Aug-98 : T : NONE : Straw removed
 : T : NORMAL, 2 NORMAL, 4 NORMAL : Straw applied and chopped.
02-Sep-98 : B : : Ploughed.
14-Sep-98 : B : : Cambridge rolled.
12-Oct-98 : B : : Rotary harrowed. Drilled Hereward, tr. Sibutol, at
 325 seeds/m² with the Accord drill.
05-Nov-98 : B : : Hardy at 15 kg.
18-Feb-99 : B : : 34.5% N at 145 kg.
25-Feb-99 : B : : tm)Platform S at 1.0 kg in 200 l.
 : B : : tm)Isoguard at 2.0 l in 200 l.
02-Apr-99 : B : : 34.5% N at 377 kg.
24-Apr-99 : B : : tm)Folicur at 0.5 l in 200 l.
 : B : : tm)Tripart Brevis at 2.0 l in 200 l.
27-May-99 : B : : tm)Folicur at 0.75 l in 200 l.
 : B : : tm)Sipcam UK Rover 500 at 1.0 l in 200 l
20-Aug-99 : B : : Combine harvested.

99/R/CS/326 GREAT KNOTT III (R)

GRAIN TONNES/HECTARE

***** Tables of means *****

STRAW	
NONE	8.83
NORMAL	7.91
2 NORMAL	7.75
4 NORMAL	7.73
Mean	8.06

*** Standard errors of differences of means ***

STRAW
1.022

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	9	1.445	17.9
GRAIN MEAN DM%	88.3		

STRAW TONNES/HECTARE

***** Tables of means *****

STRAW	
NONE	4.22
NORMAL	4.45
2 NORMAL	4.34
4 NORMAL	4.75
Mean	4.44

STRAW MEAN DM% 80.9

PLOT AREA HARVESTED 0.00310