

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 2000

[Full Table of Content](#)



00/R/CS/508 Seed Treatment Sequences and Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (2001) *00/R/CS/508 Seed Treatment Sequences and Take-all - W. Wheat* ;
Yields Of The Field Experiments 2000, pp 105 - 106 - DOI:

<https://doi.org/10.23637/ERADOC-1-55>

00/R/CS/508

SEED TREATMENT SEQUENCES AND TAKE-ALL

Object: To determine effects of a seed treatment fungicide on take-all (*Gaeumannomyces graminis*) and populations of the causal fungus - Stackyard.

Sponsors: J.F. Jenkyn, G.L. Bateman, R.J. Gutteridge.

The second year, w. wheat.

For previous year see 99/R/CS/508.

Design: 4 randomised blocks of 2 x 2 x 2 x 2.

Plot dimensions: 6.0 x 10.0.

Treatments:

- S1999** Seed dressing and year applied:
 (-99) 1999 none
 (F99) 1999 fluquinconazole
- S2000** Seed dressing and year applied:
 -00 2000 none
 F00 2000 fluquinconazole and prochloraz

Randomised for treatments as above in two subsequent years

Experimental diary:

- 02-Sep-99 : B : : Shallow (10 cm) ploughed.
- 04-Sep-99 : B : : Rolled.
- 04-Oct-99 : T : -00 : Combination drilled, Hereward, undressed, at 380 seeds/m² with the Accord drill.
- : T : F00 : Combination drilled, Hereward, tr. fluquinconazole, at 380 seeds/m² with the Accord drill.
- 06-Oct-99 : B : : Rolled.
- 12-Nov-99 : B : : tm)Lexus 50 DF at 20 g in 200 l.
- : B : : tm)Stomp 400 SC at 2.5 l in 200 l.
- : B : : tm)Toppel 10 at 250 ml in 200 l.
- 07-Mar-00 : B : : 34.5% N at 232 kg.
- 19-Apr-00 : B : : 34.5% N at 435 kg.
- 24-Apr-00 : B : : tm)Opus at 0.5 l in 100 l.
- : B : : tm)Unix at 0.5 kg in 100 l.
- : B : : tm)BASF 3C Chlormequat 720 at 2.2 l in 100 l.
- 12-May-00 : B : : tm)Ally at 30 g in 200 l.
- : B : : tm)Starane 2 at 0.5 l in 200 l.
- 22-May-00 : B : : Opus at 0.75 l in 100 l.
- 27-Jun-00 : B : : Folicur at 0.25 l in 100 l.
- 29-Jun-00 : B : : Hand rogued wild oats.
- 11-Aug-00 : B : : Combine harvested.

Previous crops: S. wheat 1998, w. wheat 1999.

NOTE: Plant and soil samples were taken in June to assess root and stem base diseases, take-all infectivity of the soil and diversity of fungal communities on roots.

00/R/CS/508

GRAIN TONNES/HECTARE

***** Tables of means *****

S2000	-00	F00	Mean
S1999			
(-99)	6.40	6.69	6.55
(F99)	6.06	6.53	6.30
Mean	6.23	6.61	6.42

*** Standard errors of differences of means ***

S1999	S2000	S1999
		S2000
0.265	0.265	0.375

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	57	1.061	16.5
GRAIN MEAN DM%	83.2		
PLOT AREA HARVESTED	0.00240		