


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 2000

[Full Table of Content](#)



00/R/WW/6 Herbicide Resistant Black-grass - W. Wheat

Rothamsted Research

Rothamsted Research (2001) *00/R/WW/6 Herbicide Resistant Black-grass - W. Wheat* ; Yields Of The Field Experiments 2000, pp 121 - 123 - DOI: <https://doi.org/10.23637/ERADOC-1-55>

00/R/WW/6

WINTER WHEAT

HERBICIDE RESISTANT BLACK-GRASS

Object: To evaluate the efficacy of different herbicide treatments on a herbicide-resistant black-grass population - Claycroft.

Sponsors: S.R. Moss.

Design: 4 randomised blocks of 15 plots.

Whole plot dimensions: 3.0 x 12.0.

Treatments:

HERBICIDE	Herbicide type, rate of active ingredient and timing (black-grass growth stage):
A	Isoproturon at 2.5 kg post emergence
B	Tri-allate at 2.25 kg pre-emergence and trifluralin at 0.96 kg with isoproturon at 1.5 kg post emergence
C	Tri-allate at 2.25 kg pre-emergence and pendimethalin at 1.32 kg with isoproturon at 1.5 kg post emergence
D	Trifluralin at 0.96 kg pre-emergence and flupyrsulfuron-methyl at 0.01 kg with pendimethalin at 1.32 kg post emergence
E	Tri-allate at 2.25 kg pre-emergence and flupyrsulfuron-methyl at 0.01 kg with pendimethalin at 1.32 kg post emergence
F	Tri-allate at 2.25 kg pre-emergence and flupyrsulfuron-methyl at 0.01 kg with trifluralin at 0.96 kg post emergence
G	Tri-allate at 2.25 kg pre-emergence and clodinafop-propargyl at 0.03 kg with trifluralin at 0.96 kg and adjuvant at two leaf stage
H	Tri-allate at 2.25 kg pre-emergence and clodinafop-propargyl at 0.03 kg with trifluralin at 0.96 kg, flupyrsulfuron at 0.01 kg and adjuvant at two leaf stage
J	Fenoxaprop-P-ethyl at 0.069 kg at two leaf stage
K	Compound K at two leaf stage
L	Compound L pre-emergence
M	Compound M pre-emergence
N	Compound N at two leaf stage with adjuvant
O	None (duplicated)

NOTE: Herbicides K, L, M and N are under commercial development, composition undisclosed.

Experimental diary:

29-Aug-99 : B : : Ploughing started.
30-Aug-99 : B : : Ploughing completed.
31-Aug-99 : B : : Rotary harrowed.
03-Sep-99 : B : : Rolled.
14-Sep-99 : B : : Disced.
16-Sep-99 : B : : Combination drilled, Consort, tr. Sibutol at 350 seeds/m² with the Nordsten drill.
21-Sep-99 : T : D : Alpha Trifluralin 48 EC at 2.0 l in 220 l.
 : T : L : Compound L at 4.0 l in 220 l.
 : T : M : Compound M at 1.0 kg in 220 l.

00/R/WW/6

Experimental diary:

22-Sep-99 : B : : Genesis at 5.0 kg.
: T : B,C,E,F,G,H : Avadex Excel 15G at 15.0 kg.
15-Oct-99 : B : : Cyperkill 10 at 250 ml in 200 l.
26-Oct-99 : T : A : Isoguard at 5.0 l in 220 l.
: T : B : tm)Isoguard at 3.0 l in 220 l.
: T : B : tm)Alpha Trifluralin 48 EC at 2.0 l in 220 l.
: T : C : tm)Isoguard at 3.0 l in 220 l.
: T : C : tm)Stomp 400 SC at 3.3 l in 220 l.
: T : D,E : tm)Lexus 50 DF at 20 g in 220 l.
: T : D,E : tm)Stomp 400 SC at 3.3 l in 220 l.
: T : F : tm)Lexus 50 DF at 20 g in 220 l.
: T : F : tm)Alpha Trifluralin 48 EC at 2.0 l in 220 l.
15-Nov-99 : T : G : tm)Hawk at 2.5 l in 220 l.
: T : G : tm)Actipron at 1.1 l in 220 l.
: T : H : tm)Hawk at 2.5 l in 220 l.
: T : H : tm)Actipron at 1.1 l in 220 l.
: T : H : tm)Lexus 50 DF at 20 g in 220 l.
: T : J : Cheetah Super at 1.25 l in 220 l.
: T : K : tm)Compound K in 220 l.
: T : N : tm)Compound N in 220 l.
: T : N : tm)Agral at 0.22 l in 220 l.
31-Mar-00 : B : : Avenge 2 at 3.3 l in 200 l.
18-Apr-00 : B : : 34.5% N at 580 kg.
29-Apr-00 : B : : Landmark at 0.5 l in 200 l.
12-May-00 : B : : tm)Ally at 30 g in 200 l.
: B : : tm)Starane 2 at 0.5 l in 200 l.
20-May-00 : B : : Landmark at 1.0 l in 100 l.
15-Aug-00 : B : : Combine harvested.

Previous crops: W. wheat 1998 and 1999.

NOTE: Black-grass populations were assessed in March and heads counted in June.

00/R/WW/6

GRAIN TONNES/HECTARE

***** Tables of means *****

HERBCIDE	
A	4.93
B	7.47
C	7.50
D	7.26
E	7.92
F	7.20
G	7.53
H	7.41
J	4.87
K	7.66
L	6.63
M	3.73
N	5.41
O	3.72
Mean	6.20

*** Standard errors of differences of means ***

HERBCIDE	
0.566	min.rep
0.490	max-min

HERBCIDE
max-min 0 v any of the remainder
min.rep Any of the remainder

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	43	0.800	12.9
GRAIN MEAN DM%	85.8		
PLOT AREA HARVESTED	0.00230		