


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 2000

[Full Table of Content](#)



---

### 00/W/WW/23 Timing of Sulphur Application on Winter Wheat

#### Rothamsted Research

Rothamsted Research (2001) *00/W/WW/23 Timing of Sulphur Application on Winter Wheat* ; Yields Of The Field Experiments 2000, pp 137 - 138 - DOI: <https://doi.org/10.23637/ERADOC-1-55>

00/W/WW/23

WINTER WHEAT

TIMING OF SULPHUR APPLICATION ON WINTER WHEAT

**Object:** To test the timing of sulphur fertilizer applications on the yield and quality of winter wheat - Woburn, Butt Close III.

**Sponsors:** M.J. Hawkesford, F.J. Zhao, M.M.A. Blake-Kalff.

**Design:** 4 randomised blocks of 5 plots.

**Plot dimensions:** 4.0 x 10.0.

**Treatments:-**

**S TIMING** Sulphur timing (30 kg S applied as gypsum):

-	None
T1	05-Mar-00
T2	06-Apr-00
T3	09-May-00
T4	07-Jun-00

**Experimental diary:**

06-Sep-99	: B :	: Ploughed and furrow pressed.
14-Sep-99	: B :	: Dutch harrowed.
	: B :	: Drilled, Hereward, tr. Sibutol, at 275 seeds/m <sup>2</sup> with the Accord drill.
12-Oct-99	: B :	: tm)Stomp 400 SC at 2.5 l in 200 l.
	: B :	: tm)Cyperkill 10 at 250 ml in 200 l.
	: B :	: tm)Isoguard at 1.0 l in 200 l.
05-Mar-00	: T : T1	: Gypsum at 171 kg.
23-Mar-00	: B :	: 34.5% N at 145 kg.
06-Apr-00	: T : T2	: Gypsum at 171 kg.
08-May-00	: B :	: tm)Vytel Copper at 0.1 l in 200 l.
	: B :	: tm)Vytel Manganese at 3.0 l in 200 l.
	: B :	: tm)Landmark at 0.5 l in 200 l.
	: B :	: tm)BASF 3C Chlormequat 720 at 2.0 l in 200 l.
09-May-00	: T : T3	: Gypsum at 171 kg.
	: B :	: 34.5% N at 377 kg.
13-May-00	: B :	: Starane 2 at 2.0 l in 200 l.
20-May-00	: B :	: Landmark at 1.0 l in 200 l.
07-Jun-00	: T : T4	: Gypsum at 171 kg.
17-Aug-00	: B :	: Combine harvested.

Previous crops: S. barley 1998, s. linseed 1999.

**NOTE:** Plant samples were taken monthly April to August for biomass, nitrogen and sulphur content.

00/W/WW/23

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

S TIMING

-	5.75
T1	7.56
T2	6.88
T3	7.40
T4	5.48

Mean	6.62
------	------

\*\*\* Standard errors of differences of means \*\*\*

S TIMING

0.402

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	12	0.568	8.6

GRAIN MEAN DM% 83.7

PLOT AREA HARVESTED 0.00192