


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 2000

[Full Table of Content](#)



00/R/LNS/1 Fungicides and Spring Linseed

Rothamsted Research

Rothamsted Research (2001) *00/R/LNS/1 Fungicides and Spring Linseed* ; Yields Of The Field Experiments 2000, pp 161 - 163 - DOI: <https://doi.org/10.23637/ERADOC-1-55>

00/R/LNS/1

LINSEED

FUNGICIDES AND SPRING LINSEED

Object: To test fungicidal spray programmes on spring linseed - Drapers.

Sponsors: B.D.L. Fitt, S.A.M. Perryman.

Design: 2 blocks of 2 plots randomly split into 3 x 4 + 2.

Plot dimensions: 3.0 x 12.0.

Treatments: All combinations of:-

Whole plots: All combinations of:-

1. VARIETY

A	Antares
J	Jupiter

Sub-plots

2. FUNGICIDE Target organism and fungicides used:

AL	<i>Alternaria</i> (Rovral Flo at 2.0 l in 220 l)
BO	<i>Botrytis</i> (Benlate Fungicide at 1.1 kg in 220 l)
A+B	<i>Alternaria</i> and <i>Botrytis</i> (Compass at 3.0 l in 220 l)

3. TIMING Fungicide timing:

P+M	Pre-flowering and mid-flowering
MF	Mid-flowering
CD	Capsule development
M+C	Mid-flowering and capsule development

Plus two extra treatments:

EXTRA

A	Antares with no fungicide (duplicated)
J	Jupiter with no fungicide (duplicated)

Experimental diary:

16-Mar-00	: B :	: Rolled. Spring-tine cultivated.
02-May-00	: T : A	: Combination drilled, Antares, tr. Prelude 20 LF at 700 seeds/m ² , with the Accord drill.
	: T : J	: Combination drilled, Jupiter, tr. Prelude 20 LF at 700 seeds/m ² , with the Accord drill.
05-May-00	: B :	: Rolled.
10-May-00	: B :	: 34.5% N at 174 kg. Hallmark with Zeon Technology at 75 ml in 200 l.
12-May-00	: B :	: Hallmark with Zeon Technology at 75 ml in 200 l.
16-May-00	: B :	: Hallmark with Zeon Technology at 75 ml in 200 l.

00/R/LNS/1

Experimental diary:

27-Jun-00 : T : P+M : Fungicides applied.
 06-Jul-00 : T : M+C,MF,P+M : Fungicides applied.
 21-Jul-00 : T : CD,M+C : Fungicides applied.
 18-Aug-00 : B : : tm)Enhance Low Foam at 400 ml in 400 l.
 18-Aug-00 : B : : tm)Reglone at 3.0 l in 400 l.
 25-Aug-00 : B : : Combine harvested.

Previous crops: Linseed 1998, w. barley 1999.

NOTE: Crop samples were taken on four occasions from June to August for disease assessments.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

FUNGICIDE	AL	BO	A+B	Mean
TIMING				
P+M	0.92	1.36	1.12	1.13
MF	1.04	1.06	1.12	1.07
CD	1.18	1.09	1.03	1.10
F+C	0.89	1.16	0.91	0.98
Mean	1.01	1.17	1.04	1.07

VARIETY	A	J	Mean
TIMING			
P+M	1.11	1.15	1.13
MF	1.16	0.98	1.07
CD	1.04	1.16	1.10
F+C	0.96	1.00	0.98
Mean	1.07	1.07	1.07

VARIETY	A	J	Mean
FUNGICIDE			
AL	1.03	0.98	1.01
BO	1.18	1.16	1.17
A+B	1.01	1.08	1.04
Mean	1.07	1.07	1.07

FUNGICIDE	AL	BO	A+B
VARIETY	A	J	A
TIMING	A	J	J
P+M	0.96	0.87	1.24
MF	1.00	1.08	1.29
CD	1.32	1.05	1.03
F+C	0.83	0.94	1.15
Mean	1.07	1.07	1.07

EXTRA	A	J	Mean
	0.83	1.05	0.94

Grand mean 1.05

00/R/LNS/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

*** Standard errors of differences of means ***

FUNGICIDE	TIMING	VARIETY*
		FUNGICIDE
0.090	0.104	0.127
VARIETY*	FUNGICIDE	VARIETY*
TIMING	TIMING	FUNGICIDE
		TIMING
0.147	0.180	0.255

* Within the same level of **VARIETY** only

S.e.d. for comparing **EXTRA** A or J with any item in **VARIETY.FUNGICIDE.TIMING** table is 0.220 within the same level of **VARIETY** only

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP.SP	28	0.255	24.2
GRAIN MEAN DM%	80.5		
PLOT AREA HARVESTED	0.00240		