Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 2000



Full Table of Content

00/R/LP/11 Yellow Lupins

Rothamsted Research

Rothamsted Research (2001) 00/R/LP/11 Yellow Lupins; Yields Of The Field Experiments 2000, pp 179 - 180 - DOI: https://doi.org/10.23637/ERADOC-1-55

00/R/LP/11

LUPINS

YELLOW LUPINS

Object: To study the effects of sowing date and seed rate upon maturity date and seed yield of yellow lupin - Highfield IV/Road Piece East.

Sponsors: I.F. Shield, H.J. Stevenson, J.E. Leach, T. Scott.

Design: 3 partially randomised blocks of 3 x 3 plots.

Whole plot dimensions: 3.0×9.0 .

Treatments: All combinations of:-

1. SOW DATE

D1	Sown	early
D2	Sown	middle
D3	Sown	late

SEED RATE Seeds per m²:

S1	70
S2	100
53	130

Experimental diary:

13-Aug-99	:	В	:		:	Ploughing started.
14-Aug-99	:	В	:		;	Ploughing completed.
24-Feb-00	:	В	:		:	PDQ at 4.0 1 in 200 1.
07-Mar-00	:	В	:		:	Spring-tine cultivated.
08-Mar-00	:	T	:	D1	:	Combination drilled, Wodjil, recleaned, with the Accord drill.
17-Mar-00	:	T	:	D1	:	Stomp 400 SC at 3.0 1 in 220 1.
28-Mar-00	:	T	:	D2	:	Combination drilled, Wodjil, recleaned, with the Accord drill.
31-Mar-00	:	T	:	D2	:	Stomp 400 SC at 3.0 1 in 220 1.
02-May-00	:	T	:	D3	:	Combination drilled, Wodjil, recleaned, with the Accord drill.
08-May-00	:	T	:	D3	:	Stomp 400 SC at 3.0 1 in 220 1.
02-Jun-00	:	В	:		:	Falcon at 1.0 l in 200 l.
30-Aug-00	:	В	:		:	tm) Enhance Low Foam at 400 ml in 400 l.
	:	В	:		:	tm) Reglone at 3.0 1 in 400 1.
12-Sep-00	:	В	:		:	Combine harvested.

Previous crops: W. barley and w. wheat 1998, w. barley 1999.

NOTES: (1) The design was intended to be a randomised block experiment but treatment combination SOW DATE D2 SEEDRATE S1 was incorrectly drilled, so plots with the correct combination were sown at the edges of the experiment. The yields were analysed using covariates to remove block effects.

00/R/LP/11

NOTES: (2) Apical dissections were made frequently between April and June. Plant densities were assessed in May, June and August. Flowering date was recorded. Branch and number of leaves were counted and plant height measured in July. Samples were taken in August from the first two sowings to assess components of yields.

GRAIN TONNES/HECTARE

**** Tables of means ****

DATE	D1	D2	D3	Mean	
S1	1.30	1.41	0.78	1.16	
S2	1.57	1.71	0.97	1.42	
S3	1.82	1.74	1.41	1.66	
Mean	1.56	1.62	1.05	1.41	

*** Standard errors of differences of means ***

SEED RATE	SOW DATE	SEED	RATE	
		SOW	DATE	
0.126	0.127	(0.220	

***** Stratum standard errors and coefficients of variation *****

Stratum d.f. s.e. cv% BLOCK.WP 16 0.266 18.9

GRAIN MEAN DM% 79.6

PLOT AREA HARVESTED 0.00206 or 0.00216 (SOW DATE D3)