

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 2000

[Full Table of Content](#)



---

### 00/R/CS/323 Cereal Sequence and Take-all - W. Wheat

#### Rothamsted Research

Rothamsted Research (2001) *00/R/CS/323 Cereal Sequence and Take-all - W. Wheat* ; Yields Of The Field Experiments 2000, pp 62 - 63 - DOI: <https://doi.org/10.23637/ERADOC-1-55>

00/R/CS/323

**CEREAL SEQUENCES AND TAKE-ALL**

**Object:** To study the effect of seed treatment on take-all (*Gaeumannomyces graminis*) in w. wheat grown after various cereal sequences - West Barnfield II.

**Sponsors:** R.J. Gutteridge, J.F. Jenkyn, G.L. Bateman.

The 13th year, w. wheat.

For previous years see 88-96,98-99/R/CS/323.

**Design:** 3 randomised blocks of 26 plots.

**Whole plot dimensions:** 3.0 x 10.0.

**Treatments:**

1. **FUNG99**                      Seed dressing in 1999:  
    (NONE)                      None  
    (FLUQUIN)                  Fluquinconazole
2. **FUNG00**                      Seed dressing in 2000:  
    NONE                        None  
    FLUQUIN                    Fluquinconazole
3. **GROUP**                      Crop sequences and levels of take-all decline:  
    I                            Old sequences with well established take-all decline  
    II                          Old sequences with less well established take-all decline  
    III                         Old sequences with take-all decline possibly not established

**Experimental diary:**

- 11-Aug-99 : B : : Ploughed.  
30-Sep-99 : B : : tm)PDQ at 3.0 l in 200 l.  
          : B : : tm)Enhance Low Foam at 100 ml in 200 l.  
11-Oct-99 : T : NONE : Combination drilled, Hereward, tr. Sibutol, at 380 seeds/m<sup>2</sup> with the Accord drill.  
          : T : FLUQUIN : Combination drilled, Hereward, tr. fluquinconazole, at 380 seeds/m<sup>2</sup> with the Accord drill.  
15-Nov-99 : B : : tm)Lexus 50 DF at 20 g in 200 l.  
          : B : : tm)Stomp 400 SC at 2.5 l in 200 l.  
          : B : : tm)Toppel 10 at 250 ml in 200 l.  
07-Mar-00 : B : : 34.5% N at 232 kg.  
27-Apr-00 : B : : 34.5% N at 348 kg.  
          : B : : tm)Opus at 0.5 l in 200 l.  
          : B : : tm)Unix at 0.5 kg in 200 l.  
          : B : : tm)BASF 3C Chlormequat 720 at 2.0 l in 200 l.  
22-May-00 : B : : Opus at 0.75 l in 100 l.  
29-Jun-00 : B : : Hand rogued wild oats.  
11-Aug-00 : B : : Combine harvested.

Previous crops: W. wheat 1998 and 1999.

**NOTE:** All plots were sampled in June to assess root and stem base diseases.

00/R/CS/323

**GRAIN TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

<b>FUNG99 GROUP</b>	(NONE)	(FLUQUIN)	Mean
I	9.23	8.90	9.06
II	9.48	9.02	9.25
III	9.04	8.65	8.84
Mean	9.25	8.85	9.05

<b>FUNG00 GROUP</b>	NONE	FLUQUIN	Mean
I	9.14	8.99	9.06
II	9.10	9.39	9.25
III	8.64	9.05	8.84
Mean	8.96	9.14	9.05

<b>FUNG00 FUNG99</b>	NONE	FLUQUIN	Mean
(NONE)	9.10	9.40	9.25
(FLUQUIN)	8.82	8.89	8.85
Mean	8.96	9.14	9.05

<b>GROUP</b>	<b>FUNG00 FUNG99</b>	NONE	FLUQUIN
I	(NONE)	9.20	9.26
	(FLUQUIN)	9.08	8.72
II	(NONE)	9.42	9.53
	(FLUQUIN)	8.78	9.26
III	(NONE)	8.67	9.41
	(FLUQUIN)	8.61	8.68

\*\*\* Standard errors of differences of means \*\*\*

<b>GROUP FUNG99</b>	<b>FUNG99</b>	<b>FUNG00</b>	<b>GROUP FUNG99</b>
0.142	0.116	0.116	0.200
<b>GROUP FUNG00</b>	<b>FUNG99 FUNG00</b>	<b>GROUP FUNG99 FUNG00</b>	
0.200	0.163	0.283	

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	58	0.490	5.4
GRAIN MEAN DM%	86.5		
PLOT AREA HARVESTED	0.00237		