

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 2000

[Full Table of Content](#)



00/R/CS/323 Cereal Sequence and Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (2001) *00/R/CS/323 Cereal Sequence and Take-all - W. Wheat* ; Yields Of The Field Experiments 2000, pp 62 - 63 - DOI: <https://doi.org/10.23637/ERADOC-1-55>

00/R/CS/323

CEREAL SEQUENCES AND TAKE-ALL

Object: To study the effect of seed treatment on take-all (*Gaeumannomyces graminis*) in w. wheat grown after various cereal sequences - West Barnfield II.

Sponsors: R.J. Gutteridge, J.F. Jenkyn, G.L. Bateman.

The 13th year, w. wheat.

For previous years see 88-96,98-99/R/CS/323.

Design: 3 randomised blocks of 26 plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments:

1. **FUNG99** Seed dressing in 1999:
 (NONE) None
 (FLUQUIN) Fluquinconazole
2. **FUNG00** Seed dressing in 2000:
 NONE None
 FLUQUIN Fluquinconazole
3. **GROUP** Crop sequences and levels of take-all decline:
 I Old sequences with well established take-all decline
 II Old sequences with less well established take-all decline
 III Old sequences with take-all decline possibly not
 established

Experimental diary:

- 11-Aug-99 : B : : Ploughed.
30-Sep-99 : B : : tm)PDQ at 3.0 l in 200 l.
 : B : : tm)Enhance Low Foam at 100 ml in 200 l.
11-Oct-99 : T : NONE : Combination drilled, Hereward, tr. Sibutol, at 380
 seeds/m² with the Accord drill.
 : T : FLUQUIN : Combination drilled, Hereward, tr. fluquinconazole,
 at 380 seeds/m² with the Accord drill.
15-Nov-99 : B : : tm)Lexus 50 DF at 20 g in 200 l.
 : B : : tm)Stomp 400 SC at 2.5 l in 200 l.
 : B : : tm)Toppel 10 at 250 ml in 200 l.
07-Mar-00 : B : : 34.5% N at 232 kg.
27-Apr-00 : B : : 34.5% N at 348 kg.
 : B : : tm)Opus at 0.5 l in 200 l.
 : B : : tm)Unix at 0.5 kg in 200 l.
 : B : : tm)BASF 3C Chlormequat 720 at 2.0 l in 200 l.
22-May-00 : B : : Opus at 0.75 l in 100 l.
29-Jun-00 : B : : Hand rogued wild oats.
11-Aug-00 : B : : Combine harvested.

Previous crops: W. wheat 1998 and 1999.

NOTE: All plots were sampled in June to assess root and stem base diseases.

00/R/CS/323

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNG99 GROUP	(NONE)	(FLUQUIN)	Mean
I	9.23	8.90	9.06
II	9.48	9.02	9.25
III	9.04	8.65	8.84
Mean	9.25	8.85	9.05

FUNG00 GROUP	NONE	FLUQUIN	Mean
I	9.14	8.99	9.06
II	9.10	9.39	9.25
III	8.64	9.05	8.84
Mean	8.96	9.14	9.05

FUNG00 FUNG99	NONE	FLUQUIN	Mean
(NONE)	9.10	9.40	9.25
(FLUQUIN)	8.82	8.89	8.85
Mean	8.96	9.14	9.05

GROUP	FUNG00 FUNG99	NONE	FLUQUIN
I	(NONE)	9.20	9.26
	(FLUQUIN)	9.08	8.72
II	(NONE)	9.42	9.53
	(FLUQUIN)	8.78	9.26
III	(NONE)	8.67	9.41
	(FLUQUIN)	8.61	8.68

*** Standard errors of differences of means ***

GROUP	FUNG99	FUNG00	GROUP FUNG99
0.142	0.116	0.116	0.200
GROUP FUNG00	FUNG99 FUNG00	GROUP FUNG99 FUNG00	
0.200	0.163	0.283	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	58	0.490	5.4
GRAIN MEAN DM%	86.5		
PLOT AREA HARVESTED	0.00237		