Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 2000



Full Table of Content

Fertilizers

Rothamsted Research

Rothamsted Research (2001) *Fertilizers*; Yields Of The Field Experiments 2000, pp 6 - 7 - **DOI:** https://doi.org/10.23637/ERADOC-1-55

CONVENTIONS 2000

For each experiment current treatments are shown with the factor and level names which are used in the tables.

For each experiment, other than annuals, references are given to previous years. These refer to the '(Numerical)(Results)' previous editions of 'Yields of the Field Experiments'.

For the classical and some long-term experiments reference is made to 'Details' - separate publications, giving full descriptions of treatments until 1977 & 1973, with full titles 'Details of the Classical and Long Term Experiments up to 1977' and 'Details of the Classical and Long Term Experiments up to 1973'.

The following conventions are observed unless otherwise stated.

All areas are in hectares. All plot dimensions are in metres.

All rates of application of fertilizers, sprays etc. are per hectare.

All yields are per hectare.

For any other crop, details of abbreviations are given as necessary.

Fertilizers

27% N or 34.5% N means nitrogen as ammonium nitrate

46% N means nitrogen as urea

Ashlade Nu Trace 5% magnesium and 1% copper

Epsom salts MgSO_{4.7H2}O 10% magnesium and 13%

sulphur

Fishmeal approximately 6.5% nitrogen

YM Farmyard manure (from bullocks)

Gypsum 17.5% sulphur

Kieserite MgSO₄H₂O 17.7% magnesium and 23.3%

sulphur

Manganese sulphate Mn₂(SO₄), 27% manganese and 24%

sulphur

Marshland Liquid Manganese Complex 150 g/l manganese, 7.5 g/l magnesium oxide (4.5 g/l Mg) and 223.6 g/l

sulphur trioxide (89.4 g/l S)

Muriate of potash 60% K,O

Nitrate of soda NaNO, 16% nitrogen and 27% sodium

Phosyn Manganese 150 g/l manganese

Profol 500 500 g/l manganese

Profol Copper 500 500 g/l copper

Profol RM 5% boron, 7% manganese, 0.4% molybdenum, 13.3% magnesium oxide (8% Mg) and 36.3% sulphur trioxide

(14.5% S)

Fertilizers (continued)

Resistim 10.9% w/w potassium and 6.3% w/w

phosphorus combined with natural

betaines

Rhodoman A seed dressing containing manganese

Silicate of soda Na,SiO, 37% sodium and 23% silica

Sulphan 30% nitrogen and 7.6% sulphur

Sulphur Gold 30% nitrogen and 7.6% sulphur

Sulphate of ammonia (NH₄)₂SO₄ 21% nitrogen 24% sulphur

Sulphate of potash K,SO, 50% K,O and 18.4% sulphur

Tiger 90 90% sulphur

Thiovit 80% sulphur

Triple superphosphate 47% P2O5

Vytel Manganese 6.4% manganese

Compound fertilizers are indicated as - (20:10:10) = (20% N, 10% P₂O₅, 10% K₂O),granular unless otherwise stated.

Cereal straw is removed unless otherwise stated.

In the experimental diary;

T: Refers to treatments applied to part of the experiment.

B: Refers to basal operations and applications to the whole experiment.

GS: Growth stage.

tm): Tank mix; two or more products applied together.

tr.: means seed dressing

Machinery definitions as used in the diary.

Accord Pneumatic drill with Suffolk coulters 12.5 cm apart.

Drill with rigid times 11.5 cm apart. Carrier Combine drilled Drill mounted behind a rotary harrow.

Dutch harrow Rigid tine harrow

Fiona Drill with Suffolk coulters 12 cm apart

Flexitine Heavy spring-tine cultivator. Hege Drill with coulters 14 cm apart

Nodet Gougis Pneumatic precision drill with variable spacing.

Drill with Suffolk coulters 12 cm apart. Nordsten Oyjord Drill with Suffolk coulters 14.2 cm apart.

Rotaspike

Spiked rotary cultivator Rotary spiked cultivator for forming potato ridges Rotaridger Deep tine cultivator with vibrating tines 60 cm apart Shakerator

and 45 cm deep. Subsoiler

Deep tine cultivator with vibrating tines 60 cm apart

and 45 cm deep

Thistlebar Shallow cultivator used to weed fallows