

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 2002

[Full Table of Content](#)

Yields of the  
Classical  
and other  
Long-term Experiments  
2002

IACR - Rothamsted

---

## 02/W/CS/478 - Woburn Continuous Maize

### Rothamsted Research

Rothamsted Research (2003) *02/W/CS/478 - Woburn Continuous Maize* ; Yields Of The Field Experiments 2002, pp 53 - 54 - DOI: <https://doi.org/10.23637/ERADOC-1-259>

02/W/CS/478

CONTINUOUS MAIZE

**Object:** To monitor the fate of organic carbon in the soil organic matter - Woburn, Stackyard A I.

**Sponsors:** P.R. Poulton, J. Gaunt.

The sixth year, forage maize and s. barley.

For previous years see 97-01/W/CS/478.

**Design:** 3 randomised blocks of 6 plots.

**Plot dimensions:** 9.0 x 25.0.

**Treatments:**

CROP	Crop and straw treatments:
M	Continuous maize, stubble incorporated
(M)B	S. barley after five years maize, stubble incorporated
MT	Maize, stubble plus 10 t maize tops incorporated
(B)M	Maize after three years of spring barley, straw removed
BT	Continuous spring barley, straw removed plus 10 t maize tops incorporated
B	Continuous spring barley, straw removed

**Experimental diary:**

31-Oct-01 : T : BT, MT: Maize tops applied at 225 kg per plot.  
05-Nov-01 : B : : Sulphate of potash at 217 kg. Triple superphosphate at 171 kg. Ploughed.  
23-Mar-02 : B : : Sting ECO at 4.0 l in 200 l.  
09-Apr-02 : B : : Potassium sulphate at 217 kg. Rotary harrowed.  
10-Apr-02 : T : (M)B, BT, B: Drilled, Optic, tr. Raxil S, at 400 seeds/m<sup>2</sup> with the Accord drill.  
11-Apr-02 : B : : Rolled.  
12-Apr-02 : T : (M)B, BT, B: Stomp 400 SC at 2.0 l in 200 l.  
08-May-02 : B : : 33.5% N at 287 kg.  
09-May-02 : T : (B)M, MT, M: Rotary harrowed. Drilled, Hudson, tr. Mesuroil, at 10 seeds/m<sup>2</sup> with the Nodet drill.  
13-Jun-02 : B : (M)B, BT, B: tm)Copper 500 at 1.0 l in 200 l.  
          : B : : tm)Landmark at 0.75 l in 200 l.  
20-Jun-02 : B : (B)M, MT, M: tm)Gesaprim at 3.0 l in 220 l.  
          : B : : tm)Cropoil at 5.0 l in 220 l.  
29-Aug-02 : T : (M)B, BT, B: Combine harvested plots for yield. Swathed straw.  
30-Aug-02 : B : (M)B, BT, B: Combine harvested all remaining barley. Swathed straw.  
31-Aug-02 : B : (M)B, BT, B: Baled and removed straw.  
12-Sep-02 : T : (B)M, MT, M: Cut discards. Cut sample areas, weighed, and sampled.

**NOTE:** Forage maize and barley grain samples were taken for N analysis.

02/W/CS/478 MAIZE

WHOLE CROP (AT 100% DRY MATTER) TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

Treatment	
M	9.08
MT	7.41
(B)M	7.87
Mean	8.12

\*\*\* Standard errors of differences of means \*\*\*

Treatment
1.465

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
Blocks.Plots	4	1.794	22.1

GRAIN MEAN DM% 28.0

PLOT AREA HARVESTED 0.00108

S. BARLEY

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

Treatment	
(M)B	5.35
BT	4.62
B	4.19
Mean	4.72

\*\*\* Standard errors of differences of means \*\*\*

Treatment
0.068

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
Blocks.Plots	4	0.084	1.8

GRAIN MEAN DM% 84.0

PLOT AREA HARVESTED 0.00525