

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 2002

[Full Table of Content](#)

Yields of the
Classical
and other
Long-term Experiments
2002

IACR - Rothamsted

02/R/BK/1 - Broadbalk

Rothamsted Research

Rothamsted Research (2003) *02/R/BK/1 - Broadbalk* ; Yields Of The Field Experiments 2002, pp 5 - 12 - DOI: <https://doi.org/10.23637/ERADOC-1-259>

02/R/BK/1

BROADBALK

Object: To study the effects of organic manures and inorganic fertilisers on continuous w. wheat. From 1968 two three-year rotations were included: potatoes, beans, w. wheat and fallow, w. wheat, w. wheat. In 1979 the first rotation was changed to fallow, potatoes, w. wheat. In 1980 the second rotation reverted to continuous w. wheat. Since 1985 part of the second rotation was added to the first to extend the rotation to fallow, potatoes, w. wheat, w. wheat, w. wheat, in 1996 the fallow was replaced by w. oats and potatoes replaced by maize in 1997.

The 159th year, w. wheat, w. oats and forage maize.

For previous years see 'Details' 1967 and 1973, Station Report for 1966, pp. 229-231; Station Report for 1968, Part 2; Station Report for 1982, Part 2, pp. 5-44 and 74-01/BK/1.

Areas harvested:

Wheat:	Section	
	0	0.00320
	1	0.00589
	4,5,6 and 7	0.00487
	8,9	0.00512
Oats:	2	0.00487
Maize:	3	0.00162

Treatments:

In 2001 a number of the treatments were changed. The treatments are now: -

Whole plots

PLOT	Fertilizers and organic manures	
	Treatments	
	Plot	from 2001
01 (FYM) N4	01	N4
21 FYMN2	2.1	FYM N2
22 FYM	2.2	FYM
03 N11	03	None
05 (P) KMg	05	(P) K Mg
06 N1 (P) KMg	06	N1 (P) K Mg
07 N2 (P) KMg	07	N2 (P) K Mg
08 N3 (P) KMg	08	N3 (P) K Mg
09 N4 (P) KMg	09	N4 (P) K Mg
10 N4	10	N4
11 N4 PMg	11	N4 P Mg
12 N1+3+1 (P) K2Mg2	12	N1+3+1 (P) K2 Mg2
13 N4 PK	13	N4 P K
14 N4 PK* (Mg*)	14	N4 P K* (Mg*)
15 N5 (P) KMg	15	N5 (P) K Mg
16 N6 (P) KMg	16	N6 (P) K Mg
17 N1+4+1 PKMg	17	N1+4+1 P K Mg
18 N1+2+1 PKMg	18	N1+2+1 P K Mg
19 N1+1+1 KMg	19	N1+1+1 K Mg
20 N4 KMg	20	N4 K Mg

02/R/BK/1

W. oats; Nitrogen and farmyard manure were not applied.

N1,N2,N3,N4,N5,N6: 48, 96, 144, 192, 240, 288 kg N as 33.5% N; to be applied at the same time as the second dressings in the split nitrogen plots for wheat and to the seedbed for forage maize.

Split N to wheat

N1+1+1, 1+2+1 etc: Rates as above. Timings: first two weeks of March, GS31 or mid-April (whichever comes first) and GS37/mid-May.

Split N to forage maize

N2+1,2+2,2+3,2+4: Rates as above. Timings: to the seedbed and post-emergence.

P: 35 kg P as triple superphosphate.

(P): (none), to be reviewed in 2004/5.

K: 90 kg K as potassium sulphate.

K2: 180 kg K as potassium sulphate (plus 450 kg K autumn 2000 only).

K*: 90 kg K as potassium chloride.

Mg: 12 kg Mg as kieserite.

Mg2: 24 kg Mg as kieserite (plus 60 kg Mg, autumn 2000 only).

(Mg*): (none), to be reviewed in 2004/5.

FYM: Farmyard manure at 35 t

Previous treatment: -

Whole plots

PLOT	Plot	Fertilizers and organic manures:-		
		Treatments until 1967	Treatments from 1968	Treatments from 1985 - 2000
01DN4PK	01	-	D N2 P K	D N4 P K
21DN2	21	D	D N2	D N2
22D	22	D	D	D
030	03	None	None	None
05F	05	P K Na Mg	P K (Na) Mg	PK Mg
06N1F	06	N1 P K Na Mg	N1 P K (Na) Mg	N1 P K Mg
07N2F	07	N2 P K Na Mg	N2 P K (Na) Mg	N2 P K Mg
08N3F	08	N3 P K Na Mg	N3 P K (Na) Mg	N3 P K Mg
09N4F	09	N*1 P K Na Mg	N4 P K (Na) Mg	N4 P K Mg
10N2	10	N2	N2	N2
11N2P	11	N2 P	N2 P	N2 P
12N2PNA	12	N2 P Na	N2 P Na	N2 P Na
13N2PK	13	N2 P K	N2 P K	N2 P K
14N2PKMG	14	N2 P Mg	N2 P K Mg	N2 P K Mg
15N5F	15	N2 P K Na Mg	N3 P K (Na) Mg	N5 P K Mg
16N6F	16	N*2 P K Na Mg	N2 P K (Na) Mg	N6 P K Mg
17N1+3FH	17	N2 (A)	N2 2 (P K (Na) Mg)	N1+3 2 (PK Mg) (A)+
18N0+3FH	18	P K Na Mg (A)	N2 2 (P K (Na) Mg)	N0+3 2 (PK Mg) (A)+
19 (C)	19	C	C	(C) (since 1989)
20N2KMG	20	N2 K Na Mg	N2 K (Na) Mg	N2 K Mg

(A) Alternating each year

02/R/BK/1

+ This change since 1980. Treatments shown are those to w. wheat; autumn N alternates. Maize received N3 2(PK Mg) on both plots 17 and 18.

W. oats; Nitrogen and dung were not applied.

N1,N2,N3,N4,N5,N6: 48, 96, 144, 192, 240, 288 kg N as sulphate of ammonia until 1967, except N* which was nitrate of soda. All as 'Nitro-Chalk' in spring from 1968 to 1985, as 34.5% N since 1986.

N0+3; N1+3: None in autumn + 144 kg N in spring; 48 kg N in autumn + 144 kg N in spring

P: 35 kg P as triple superphosphate in 1974 and since 1988, single superphosphate in other years

K: 90 kg K as sulphate of potash

Na: 55 kg Na as sulphate of soda

(Na): 16 kg Na as sulphate of soda until 1973

Mg: 30 kg Mg annually to Plot 14 (applied at 26 kg 1990 to 2000), 35 kg Mg every third year to other plots since 1974 (applied at 30 kg in 1991, 1994, 1997 and 2000 and at 15 kg on half rate treatments). All as kieserite since 1974, previously as sulphate of magnesia annually

D: Farmyard manure at 35 t

(C): Castor meal to supply 96 kg N until 1988, none since

F: Full rate P K (Na) Mg as above H: Half rate of above

Strips of sub-plots: Until 1967 wheat alone was grown on the experiment, with some bare fallowing. From 1968, ten strips of sub-plots (sections) were started with the following cropping:-

SECTION	1/W34	9/W42	0/W49	8/W6	6/W23	5/O	3/W3	7/W1	4/M	2/W2
Section	1	9	0*	8+	6**	5	3	7	4	2
Year										
1968	W	W	W	W	F	W	W	P	W	BE
1969	W	W	W	W	W	F	W	BE	P	W
1970	W	W	W	W	W	W	F	W	BE	P
1971	W	W	W	W	F	W	W	P	W	BE
1972	W	W	W	F	W	F	W	BE	P	W
1973	W	W	W	W	W	W	F	W	BE	P
1974	W	W	W	W	F	W	W	P	W	BE
1975	W	W	W	W	W	F	W	BE	P	W
1976	W	W	W	W	W	W	F	W	BE	P
1977	W	W	W	W	F	W	W	P	W	BE
1978	W	W	W	W	W	F	W	BE	P	W
1979	W	W	W	W	W	W	F	W	P	F
1980	W	W	W	W	W	W	W	F	W	P
1981	W	W	W	F	W	W	W	P	F	W
1982	W	W	W	W	W	W	W	W	P	F
1983	W	W	W	W	W	W	W	F	W	P
1984	W	W	W	W	W	W	W	P	F	W
1985	W	W	W	W	W	F	W	W	P	W
1986	W	W	W	W	W	P	F	W	W	W
1987	W	W	W	W	W	W	P	W	W	F
1988	W	W	W	F	W	W	W	F	W	P

02/R/BK/1

SECTION	1	9	0*	8+	6**	5	3	7	4	2
Section	1	9	0*	8+	6**	5	3	7	4	2
Year										
1989	W	W	W	W	W	W	W	P	F	W
1990	W	W	W	W	W	F	W	W	P	W
1991	W	W	W	W	W	P	F	W	W	W
1992	W	W	W	W	W	W	P	W	W	F
1993	W	W	W	W	W	W	W	F	W	P
1994	W	W	W	F	W	W	W	P	F	W
1995	W	W	W	W	W	F	W	W	P	W
1996	W	W	W	W	W	P	O	W	W	W
1997	W	W	W	W	W	W	M	W	W	O
1998	W	W	W	W	W	W	W	O	W	M
1999	W	W	W	W	W	W	W	M	O	W
2000	W	W	W	W	W	O	W	W	M	W
2001	W	W	W	F	W	M	O	W	W	W
2002	W	W	W	W	W	W	M	W	W	O

W = w. wheat, O = w. oats (spring oats 2001), P = potatoes, BE = s. beans,
 F = fallow,
 M = forage maize

* Straw incorporated since autumn 1986. ** No sprays except weedkillers since 1985. + No weedkillers.

NOTES: (1) For a fuller record of treatments see 'Details' etc.
 (2) From autumn 1975 to autumn 1986, chalk was applied at 2.9 t each autumn to all plots in sets of Sections on a three-year cycle. Year 1: Sections 1,2,3. Year 2: Sections 6,7,8,9. Year 3: Sections 0,4,5. From autumn 1988 until autumn 1992 a five-year cycle was used. Year 1: Sections 1,3. Year 2: Sections 2,8. Year 3: Sections 7,9. Year 4: Sections 4,6. Year 5: Sections 0,5. None applied since autumn 1991.

Experimental diary:

All sections:

12-Sep-01 : T : FYM : FYM at 35.0 tonnes, strips 2.1 & 2.2, not oat section.
 : T : K* : Muriate of potash at 181 kg, strip 14.
 : T : P : TSP at 171 kg, strips 11, 13, 14, 17, & 18.
 : B : : Ploughing started.
 13-Sep-01 : B : : Ploughing completed.
 07-Mar-02 : T : K : Sulphate of potash at 217 kg, strips 5, 6, 7, 8, 9, 13, 15, 16, 17, 18, 19 & 20.
 : T : K2 : Sulphate of potash at 434 kg, strip 12.
 08-Mar-02 : T : MG : Kieserite at 80 kg, strips 5, 6, 7, 8, 9, 11, 15, 16, 17, 18, 19, & 20.
 : T : MG2 : Kieserite at 160 kg, strip 12.
 04-Jul-02 : B : : Rogued wild oats, 2 plants.

Cropped sections:

W. wheat

24-Aug-01 : T : : straw baled (sections 1, 4, 6, 7 & 9)
 25-Aug-01 : T : : chopped straw, section 0.
 28-Sep-01 : T : : Combination drilled, Hereward, tr. Sibutol, at 380 seeds/m² with the Accord drill.
 11-Oct-01 : T : : Avadex Excel 15 G at 15.0 kg, excluding section 8.
 02-Nov-01 : B : : tm)Lexus 50 DF at 20 g in 200 l, excluding section 8.

02/R/BK/1

Experimental diary:

Cropped sections:

W. wheat

02-Nov-01 : B : : tm)Stomp 400 SC at 3.0 l in 200 l, excluding section 8.
 15-Nov-01 : T : : Hallmark with Zeon technology at 25 ml in 200 l, not section 6.
 26-Mar-02 : T : : 1st split N applied (33 kg N over in error).
 03-Apr-02 : B : : tm)Topik at 0.125 l in 100 l, excluding section 8.
 : B : : tm)Phase II at 1.0 l in 100 l, excluding section 8.
 04-Apr-02 : B : : tm)Opus at 0.4 l in 100 l, excluding section 6.
 : B : : tm)Twist at 0.6 l in 100 l, excluding section 6.
 : B : : tm)BASF 3C chlormequat 720 at 1.25 l in 100 l, excluding section 6.
 : B : : tm)Moddus at 0.2 l in 100 l, excluding section 6.
 24-Apr-02 : T : : Main N (except plot 2.1) and 2nd split N applied (33 kg N less to split N plots to balance error).
 25-Apr-02 : T : : N to strip 2.1.
 23-May-02 : T : : 3rd split N applied.
 01-Jun-02 : T : : Opera at 1.0 l in 200 l, excluding section 6.
 14-Aug-02 : P : : Combine harvested, all discards to open up plots.
 : T : : Combine harvested plots for yield, sections 0, 1, 8, and 9.
 : P : : Swathed straw. Baled straw. Including section 0.
 15-Aug-02 : T : : Combine harvested plots for yield, completed.
 : T : : Combined side discards to allow straw weights.
 : T : : Swathed straw.
 : T : : Sampled, baled and weighed straw, sections 1 and 9.
 16-Aug-02 : T : : Sampled, baled and weighed straw, completed.
 17-Aug-02 : B : : Combine harvested all remaining wheat.
 : B : : Swathed straw. Baled straw.
 21-Aug-02 : : : Carted bales.

W. oats

24-Aug-01 : T : : straw baled, section 2
 27-Sep-01 : T : : Combination drilled, Gerald, tr. Sibutol, at 350 seeds/m² with the Accord drill.
 01-Nov-01 : B : : tm)Lexus Class at 60 g in 200 l.
 : B : : tm)Hallmark with Zeon technology at 40 ml in 200 l.
 01-Jun-02 : T : : Opera at 1.0 l in 200 l, excluding section 6.
 14-Aug-02 : P : : Combine harvested discards to allow straw weights to be taken.
 : T : : Combine harvested plots for yield.
 : T : : Swathed straw. Sampled, baled and weighed straw.
 21-Aug-02 : : : Carted bales.

Forage maize

24-Aug-01 : T : : Straw baled, section 3.
 25-Mar-02 : : : Sting ECO at 4.0 l in 200 l.
 10-Apr-02 : P : : Flexitined.
 02-May-02 : T : : Main N and 1st split N applied.
 03-May-02 : T : : Rotary harrowed.
 : T : : Drilled, Hudson, tr. Mesurol at 102,000 seeds/ha, with the Nodet Gougis drill.
 23-May-02 : T : : 2nd split N applied.
 19-Jun-02 : T : : tm)Barclay Mutiny at 0.6 l in 200 l.
 : T : : tm)Lentagran WP at 1.5 kg in 200 l.
 11-Sep-02 : T : : Cut all discard maize. Cut sample areas by hand, weighed, and sampled.
 02-Oct-02 : T : : Topped stalks ready for ploughing Countryside Stewardship Scheme margins
 10-Apr-02 : T : : Flexitined.

24-Apr-02 : P : : Drilled special headland mix at 16.0 kg with the Moore drill, 2.0 m headland strips.

02/R/BK/1

NOTE: Straw on Section 0 was baled and removed at harvest 2002 (usually incorporated) as this section will remain unploughed to test control of *Equisetum*. Samples of wheat and oat grain and straw, and forage maize were taken for chemical analysis. Unground wheat grain and straw and maize samples from selected treatments were archived.

W. WHEAT

GRAIN TONNES/HECTARE

***** Tables of means *****

SECTION PLOT	5/W1	8/W1	4/W2	7/W3	6/W25	1/W36	9/W44	0/W51
01 (FYM)N4	11.24	*	10.22	8.78	7.23	*	*	*
21FYMN2	10.46	3.53	9.47	8.05	7.11	7.66	9.26	6.87
22FYM	5.86	3.47	5.44	4.52	4.93	5.45	6.13	4.22
03Nil	1.54	2.15	0.58	0.16	0.37	0.47	0.34	0.29
05 (P) KMg	1.68	2.90	0.78	0.08	0.33	0.41	0.72	0.45
06N1 (P) KMg	3.87	2.24	2.63	0.98	1.34	1.51	1.82	1.36
07N2 (P) KMg	5.85	2.09	4.34	1.86	2.05	2.72	2.71	3.29
08N3 (P) KMg	7.37	2.55	5.82	3.11	2.70	3.08	3.30	3.55
09N4 (P) KMg	9.13	2.18	6.85	4.21	5.13	5.73	5.89	6.06
10N4	9.81	2.37	4.18	0.54	0.51	0.66	0.38	0.93
11N4PMg	8.28	2.10	7.20	3.39	2.32	4.45	3.80	4.55
12N1+3+1 (P) K2Mg2	9.97	2.05	6.25	3.08	3.94	5.05	3.85	5.18
13N4PK	8.14	2.51	5.20	3.89	4.03	5.31	5.15	4.12
14N4PK* (Mg*)	8.80	4.05	4.79	3.03	4.63	5.44	5.13	4.30
15N5 (P) KMg	10.01	2.54	6.58	4.67	4.12	4.02	6.68	4.51
16N6 (P) KMg	10.61	2.48	8.83	5.58	6.79	7.15	8.00	7.34
17N1+4+1PKMg	10.66	1.74	7.51	6.16	6.22	6.12	7.65	6.19
18N1+2+1PKMg	9.16	2.16	7.75	4.55	5.51	3.89	6.82	4.85
19N1+1+1KMg	7.47	2.38	5.81	2.90	2.25	2.54	4.32	2.14
20N4KMg	*	*	*	*	*	0.14	*	0.45

GRAIN MEAN DM% 84.2

02/R/BK/1 W.WHEAT

STRAW TONNES/HECTARE

***** Tables of means *****

SECTION PLOT	5/W1	6/W25	1/W36	9/W44
01 (FYM) N4	6.97	4.53	*	*
21 FYMN2	5.47	4.41	3.54	5.29
22 FYM	3.84	3.94	2.74	3.08
03 Nil	0.50	1.45	0.13	0.17
05 (P) KMg	0.62	1.42	0.11	0.70
06 N1 (P) KMg	1.65	0.77	0.70	1.23
07 N2 (P) KMg	2.93	0.91	1.49	1.46
08 N3 (P) KMg	3.59	1.54	1.63	2.17
09 N4 (P) KMg	4.47	2.56	2.31	3.72
10 N4	4.34	0.47	0.44	0.35
11 N4 PMg	3.33	0.76	1.90	2.23
12 N1+3+1 (P) K2Mg2	4.53	1.88	2.53	2.60
13 N4 PK	4.18	1.81	2.63	3.08
14 N4 PK* (Mg*)	4.12	1.61	2.37	2.51
15 N5 (P) KMg	5.21	2.27	2.05	3.99
16 N6 (P) KMg	5.32	3.15	2.99	5.09
17 N1+4+1 PKMg	5.38	3.05	3.67	4.80
18 N1+2+1 PKMg	3.73	2.54	1.82	4.23
19 N1+1+1 KMg	3.76	1.98	2.03	3.56
20 N4 KMg	*	*	0.09	*

STRAW MEAN DM% 87.5

W. OATS

GRAIN TONNES/HECTARE

***** Tables of means *****

PLOT	GRAIN	STRAW
01 (FYM) [N4]	6.44	3.66
21 [FYMN2]	7.63	4.29
22 [FYM]	8.29	5.82
03 Nil	2.22	0.92
05 (P) KMg	2.64	1.12
06 [N1] (P) KMg	3.48	1.46
07 [N2] (P) KMg	4.06	1.89
08 [N3] (P) KMg	4.71	2.08
09 [N4] (P) KMg	4.91	2.76
10 [N2]	5.50	3.46
11 [N2] PMg	5.85	2.82
12 [N2] (P) K2Mg2	4.39	1.89
13 [N2] PK	3.90	1.58
14 [N2] PK* (Mg*)	4.14	1.65
15 [N5] (P) KMg	3.89	1.37
16 [N6] (P) KMg	5.75	3.36
17 [N1+3] PKMg	5.44	2.32
18 [N0+3] PKMg	3.41	1.28
19 KMg	2.48	0.84

GRAIN MEAN DM% 85.9

STRAW MEAN DM% 72.9

02/R/BK/1 MAIZE

WHOLE CROP (100% DM) TONNES/HECTARE

***** Tables of means *****

PLOT	WHOLE CROP
01 (FYM) N4	10.65
21 FYMN2	7.68
22 FYM	8.41
03 Nil	2.53
05 (P) KMg	1.94
06 N1 (P) KMg	8.81
07 N2 (P) KMg	10.70
08 N3 (P) KMg	12.41
09 N4 (P) KMg	13.60
10 N4	0.67
11 N4 PMg	4.60
12 N2+3 (P) K2Mg2	12.05
13 N4 PK	12.37
14 N4 PK* (Mg*)	13.45
15 N5 (P) KMg	12.77
16 N6 (P) KMg	12.73
17 N2+4 PKMg	11.94
18 N2+2 PKMg	11.23
19 N2+1 KMg	6.15

CROP MEAN DM% 25.3